



Daniel Puppe | Manfred Wanner | Michael Sommer

Data on euglyphid testate amoeba densities, corresponding protozoic silicon pools, and selected soil parameters of initial and forested biogeosystems

Suggested citation referring to the original publication:

Data in Brief 21(2018), 1697–1703

DOI <https://doi.org/10.1016/j.dib.2018.10.164>

ISSN (online) 2352-3409

Postprint archived at the Institutional Repository of the Potsdam University in:

Postprints der Universität Potsdam

Mathematisch-Naturwissenschaftliche Reihe ; 1039

ISSN 1866-8372

<https://nbn-resolving.org/urn:nbn:de:kobv:517-opus4-471160>

DOI <https://doi.org/10.25932/publishup-47116>



ELSEVIER

Contents lists available at ScienceDirect

Data in Brief

journal homepage: www.elsevier.com/locate/dib



Data Article

Data on euglyphid testate amoeba densities, corresponding protozoic silicon pools, and selected soil parameters of initial and forested biogeosystems



Daniel Puppe^{a,*}, Manfred Wanner^b, Michael Sommer^{a,c}

^a Leibniz Centre for Agricultural Landscape Research (ZALF), Eberswalder Str. 84, 15374 Müncheberg, Germany

^b Brandenburg University of Technology Cottbus-Senftenberg, Department Ecology, 03013 Cottbus, Germany

^c University of Potsdam, Institute of Earth and Environmental Sciences, 14476 Potsdam, Germany

ARTICLE INFO

Article history:

Received 20 April 2018

Received in revised form

4 October 2018

Accepted 30 October 2018

Available online 10 November 2018

Keywords:

Silicon cycling

Biogenic silica

Terrestrial biogeosystems

Biosilicification

Euglyphida

ABSTRACT

The dataset in the present article provides information on protozoic silicon (Si) pools represented by euglyphid testate amoebae (TA) in soils of initial and forested biogeosystems. Protozoic Si pools were calculated from densities of euglyphid TA shells and corresponding Si contents. The article also includes data on potential annual biosilicification rates of euglyphid TA at the examined sites. Furthermore, data on selected soil parameters (e.g., readily-available Si, soil pH) and site characteristics (e.g., soil groups, climate data) can be found. The data might be interesting for researchers focusing on biological processes in Si cycling in general and euglyphid TA and corresponding protozoic Si pools in particular.

© 2018 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

* Corresponding author.

E-mail address: daniel.puppe@zalf.de (D. Puppe).

Specifications table

Subject area	<i>Biology</i>
More specific subject area	<i>Biogeochemistry</i>
Type of data	<i>Tables</i>
How data was acquired	<i>Soil analyses (for details see Section 2.2.) and microscopical examinations (for details see Section 2.3.)</i>
Data format	<i>Analyzed data</i>
Experimental factors	<i>Soil parameters, euglyphid testate amoeba densities, and corresponding protozoic silicon pools</i>
Experimental features	<i>Analyses of initial (chronosequence) and forest sites</i>
Data source location	<i>Germany, for latitudes & longitudes and further site details see Table 1.</i>
Data accessibility	<i>All data are presented within the paper.</i>

Value of the data

- The provided dataset is useful for comparison with the results of other authors regarding, e.g., ecological (euglyphid testate amoeba densities) and biogeochemical (protozoic silicon pools, annual biosilicification) issues in different global ecosystems.
- Data on protozoic silicon pools and corresponding annual biosilicification rates might emphasize the need for detailed investigations of silicon (re-)cycling in unicellular organisms in general and testate amoeba in particular (e.g., qualitative characterization of biogenic silicon, isotope analysis).
- Together with other datasets the presented data allow meta-analyses to examine significant controls (steady state soils) and dynamics (initial soils) of euglyphid testate amoeba densities and related amoebal biosilicification processes in soils in detail.
- In combination with data of other authors the presented data can be used for modelling to assess the role of euglyphid testate amoebae (compared to other organisms that synthesize biogenic silicon, e.g., plants or diatoms) for silicon cycling in soils and corresponding silicon fluxes from terrestrial to aquatic ecosystems.

1. Data

Diverse unicellular and multicellular organisms are able to synthesize structures of amorphous silica ($\text{SiO}_2 \cdot n\text{H}_2\text{O}$) [1]. In soils these structures represent different biogenic silicon (BSi) pools depending on their origin [2,3]. In general, BSi plays an important role in the global cycling of Si [4–6]. However, while research has been focused on plantal Si and corresponding phytogenic Si pools since decades [e.g., 7–9], far less is known about other BSi pools, e.g., protozoic Si pools represented by euglyphid testate amoeba (TA) shells [3,10–15]. Euglyphid TA represent a monophyletic clade (Euglyphida) of unicellular soil protists with a self-secreted siliceous test (shell) and a worldwide distribution [16].

The presented data were the basis for analyses of protozoic Si pools in initial [12] and forested biogeosystems [3]. The dataset in the present article provides information on i) site characteristics, geographic positions, and climate data of the initial and forested biogeosystems (Table 1), ii) analyzed soil parameters (Table S1), iii) densities of euglyphid TA shells in soils (Table S2), and iv) corresponding protozoic Si pools as well as annual biosilicification rates of euglyphid TA at these sites (Table S3).

Table 1
Site characteristics, geographic positions, and climate data of the examined initial and forested biogeosystems. Data on precipitation and temperature represent annual averages for the period 1981–2010 (German Meteorological Service).

Site	Lithology	Soil group (WRB)	Humus form	Ecosystem	Coverage (%)			Geographic position		Altitude (m a.s.l.)	Precipitation (mm)	Temperature (°C)
					Tree	Shrub	Herb	Latitude	Longitude			
CC	Spoil of Quaternary sands	Arenosol	–	Initial (artificial catchment)	–	–	–	51°36'18" N	14°15'58" E	135	568	9.6
NL	Spoil of Quaternary sands	Arenosol	–	Initial (artificial catchment)	–	–	–	51°35'50" N	14°17'22" E	132	568	9.6
AB	Calcareous glacial till	Stagnosol	Mull	Forest (beech, oak)	68	3	70	54°12'23" N	13°02'21" E	34	654	8.6
EG	Dolomitic limestone	Cambisol	Mull	Forest (beech)	65	0	78	48°20'19" N	09°25'42" E	786	808	9.5
HE	Basalt	Vertisol	Mull	Forest (beech, ash, spruce, maple)	65	3	60	47°47'53" N	08°45'08" E	703	820	9.2
MR	Calcareous loess	Luvisol	Mull	Forest (beech)	85	35	50	49°07'13" N	08°40'38" E	217	776	10.0
PP	Calcareous loess	Stagnosol	Mull	Forest (pine, beech)	70	0	2	48°51'17" N	08°46'14" E	473	889	9.5
ZE	Kaolinitic claystone	Stagnosol	Mull	Forest (hornbeam, oak)	85	10	60	48°38'37" N	09°33'13" E	390	1090	9.9
HS	Siliceous sandstone	Podzol	Moder	Forest (pine, spruce)	73	18	63	48°34'21" N	08°20'60" E	832	1607	8.1
RO	Calcareous glacial till	Luvisol	Moder	Forest (beech, oak)	70	5	5	53°30'38" N	13°21'43" E	71	530	8.7
SL	Eolian sands (dune)	Arenosol	Moder	Forest (pine, beech)	80	0	5	49°20'51" N	08°38'14" E	123	861	11.0
HK	Siliceous sandstone	Planosol	Peat	Forest (spruce, pine)	65	15	60	48°34'13" N	08°20'53" E	907	1607	8.1

2. Experimental design, materials and methods

2.1. Site description and sampling scheme

2.1.1. Initial sites

The artificial catchments 'Chicken Creek' (CC) and 'Neuer Lugteich' (NL) are part of a post-mining landscape located in the active mining area 'Welzow-South' (lignite open-cast mining, 150 km south-east of Berlin) in the state of Brandenburg, Germany. Climate is characterized by an average air temperature of 9.6 °C with an annual precipitation of 568 mm comprising data from 1981 to 2010.




Fig. 1. Photographs of initial (a-c) and some selected forest sites (d-f). A) CC (3-year-old), b) CC (5-year-old), c) NL (10-year-old), d) EG, e) HS, and f) RO. For site details see [Table 1](#) and [Section 2.1](#).

The construction of CC was completed in 2005 (time zero). In 2008 a small area in the west of the catchment was again restored to time zero (removal of the upper 20 cm of soil) for additional experimental plots. Construction of NL was finished in 2001 (time zero). Soils classify as Protic Arenosol (Calcaric, Transportic) or Haplic Arenosol (Hyperochric, Transportic) depending on site age [17]. Detailed information on site construction of CC and NL can be found in Gerwin et al. [18] and Kendzia et al. [19], respectively. All samples were taken from Quaternary substrate at 3-, 5- (CC), and 10-year-old (NL) spots representing a chronosequence (Fig. 1a-c). Samples (20 cm × 20 cm × 5 cm; subdivided in two compartments: 0–2.5 and 2.5–5 cm depth) were taken at randomly chosen spots within an area of approx. 25 m². Vegetated (cov) and uncovered (unc) spots were sampled in four field replicates each to analyze possible impacts of vegetation (3 cov: *Tussilago farfara* and *Trifolium arvense*; 5 cov: *Corynephorus canescens* and *T. arvense*) on protozoic Si pools. At NL almost the whole surface was vegetated with biogenic crusts, Poales, and several shrubs, which is why only vegetated spots (10 cov) were sampled. Samples were taken in May 2010 (CC: 5 unc, 5 cov), May 2011 (NL: 10 cov), and August 2011 (CC: 3 unc, 3 cov).

2.1.2. Forest sites

Ten non-eroded forest sites showing huge differences in climate, parent material, and pedogenesis were selected. Mean annual precipitation rates range from 530 to 1600 mm, mean annual air temperatures from 8 to 11 °C. Soils comprise (i) a sandy Arenosol developed from eolian sands (SL, dune), a Podzol and a Planosol from siliceous sandstones (HS, HK) very low in weatherable minerals (< 10% feldspars, mica), (ii) silty to loamy Luvisols and Stagnosols from calcareous, illitic loess (MR, PP) and sandy to loamy Luvisols and Stagnosols from glacial till (RO, AB), both parent materials with intermediate contents of weatherable minerals (feldspars, mica), (iii) a clayey Cambisol from dolomitic limestone (EG), (iv) a clayey Stagnosol from kaolinitic claystone (ZE), and finally (v) a clayey, smectitic Vertisol from basalt (HE) very high in weatherable minerals, like augite and plagioclase. The forest stands are old and are assumed to be in steady state in terms of TA dynamics at a decadal time scale (photographs of some selected forest sites can be found in Fig. 1d-f). Soil samples were taken in four field replicates ($n = 4$) at all sample sites except HK ($n = 3$). The field replicates (20 cm × 20 cm each) were placed randomly within an area of approx. 100 m². Per field replicate samples were taken in the upper 5 cm (incl. organic layers except for fresh litter) differentiating between two superimposed soil compartments about the same size (ideally 20 cm × 20 cm × 2.5 cm each) and transferred to plastic bags. Sampling took place in spring 2010 within six weeks (April 26th–June 6th).

2.2. Soil parameters

Bulk densities (BD, g cm⁻³) were calculated by dividing weights of oven-dried (105 °C) aliquots of soil samples by corresponding volumes. Remaining soil samples were air dried and sieved (2 mm) to separate fine-earth (< 2 mm) from skeleton content (> 2 mm). For soil analyses only fine-earth was used.

2.2.1. Soil pH, carbon, and nitrogen

Soil pH was measured using a glass electrode in a 0.01 M CaCl₂ solution with a soil-to-solution ratio of one-to-five. For total carbon as well as nitrogen analyses (C_t and N_t) fine-earth samples were finely powdered in a disc mill. Subsequently, C_t and N_t were determined by dry combustion using an elemental analyzer (CNS TruSpec, Leco Instruments). Total inorganic carbon (TIC) was measured with a multiphase analyzer (RC 612, Leco Instruments). Soil organic carbon (SOC) concentrations were calculated by subtraction (C_t-TIC) and C:N ratios were calculated by division (SOC:N_t). Soil C and N analyses were performed at the minimum of two lab repetitions per sample.

2.2.2. Readily-available silicon

For extraction of the calcium chloride (CaCl₂) soluble, so-called readily- or plant-available Si fraction (Si_{CaCl2}), 2 g of soil was mixed with 20 ml of a 0.01 M CaCl₂ solution per sample and continuously shaken for 16 h using a lab roller mixer [20]. This Si fraction was extracted to characterize the Si supply for shell synthesis of euglyphid TA in soils. Subsequent to extraction, the extracts were

centrifuged (4000 rpm, 30 min), filtered using 0.45 μm polyamide membrane filters, and Si concentrations were determined by ICP-OES (iCAP 6300 Duo, Thermo Scientific). Complete extraction work was done using plastic equipment only and results represent arithmetic means of three lab repetitions per sample.

2.2.3. Data conversion and calculation steps

All results except for pH were converted to an oven-dry basis (105 °C). Fine-earth mass (FEM in kg m^{-2}) was calculated considering bulk density, thickness and skeleton content (wt%). Total FEM (FEM_t) of the upper 5 cm was calculated as the sum of FEM of superimposed compartments. For the upper 5 cm of soil pH was averaged as follows: Per compartment pH was multiplied with the corresponding FEM, divided by FEM_t and subsequently these results were summed up. Mass densities (g m^{-2}) of SOC, N_t, and Si_{CaCl2} were calculated compartment-wise by multiplying FEM with element concentrations (g kg^{-1}). Finally, the results of superimposed compartments were summed up for the upper 5 cm of soil.

2.3. Euglyphid testate amoeba densities, protozoic silicon pools, and annual biosilicification

Soil samples in the plastic bags were homogenized by gentle manual mixing and subsequently 2 g of fresh soil was taken per sample for TA analyses and stored in 8 ml of formalin (4%). Soil suspensions received from serial dilution (1000–31.25 mg soil in 8 ml of water each) were stained with aniline blue. TA were enumerated using an inverted microscope (OPTIKA XDS-2, magnifications of 200 \times and 400 \times) differentiating between full (living incl. encysted individuals, stained) and empty shells (unstained) of the order Euglyphida. TA densities (shells cm^{-2}) were calculated considering TA shell numbers (g^{-1} dry weight), bulk density (g cm^{-3}), and thickness (cm) per soil compartment. TA densities of the upper 5 cm were calculated by summing up the corresponding TA densities of superimposed soil compartments.

For calculation of protozoic Si pools we differentiated between different TA taxa of the order Euglyphida with known silica contents per shell as published by Aoki et al. [10]. Summarizing these data we calculated Si contents (pg shell^{-1} , in parentheses listed below) by simple multiplication (SiO_2 content $\times 28/60 = \text{Si content}$) for 9 TA taxa: *Assulina muscorum* (750), *Corythion dubium* (580), *Euglypha* spec. (720), *Euglypha rotunda/laevis* type (420), *Euglypha strigosa* type (1420), *Tracheleuglypha dentata* (750), *Trinema complanatum* (500), *Trinema enchelys* (770), and *Trinema lineare* (360). Indistinctly euglyphid shells or other silica platelet synthesizing TA taxa (e.g., *Valkanovia elegans*) were recorded as 'euglyphid TA' (700 pg Si per shell, mean of the Si content per shell of the 9 taxa above).

Protozoic Si pools (BSi_{TA}; mg m^{-2}) were calculated per soil compartment using the following formula:

$$\text{BSi}_{\text{TA}} = \sum_{i=1}^n (N_i \times \text{Si}_i \times \rho_b \times t \times 10^{-5}) \quad (1)$$

where N_i is the number of euglyphid TA shells (g^{-1} dry weight), Si_i is the corresponding Si content (pg shell^{-1} ; given in parentheses listed above), ρ_b is the bulk density (g cm^{-3}), and t is the thickness (cm) of the corresponding soil compartment. In contrast, for estimation of annual biosilicification only living euglyphid TA (g^{-1} dry weight) were considered for N_i in Eq. (1) due to their ability of reproduction. After calculation steps as described in Eq. (1) results were multiplied with 13 and 90 (potential TA generations per year, see Foissner [21]) for minimal and maximal annual biosilicification rates, respectively. For calculation of protozoic Si pools and euglyphid TA biosilicification rates of the upper 5 cm of soil the results of superimposed soil compartments were added up.

Acknowledgements

DP is funded by the Deutsche Forschungsgemeinschaft (DFG) under Grant PU 626/2-1 (*Biogenic silicon in agricultural landscapes (BiSiAL) – Quantification, qualitative characterization, and importance for Si balances of agricultural biogeosystems*). DP also was partly financed by the DFG project *Spatiotemporal dynamics of biogenic Si pools in initial soils and their relevance for desilication* of MS and MW (Grant SO 302/7-1). The publication of this article was funded by the Open Access Fund of the Leibniz Association.

The authors would like to thank David M. Wilkinson and two anonymous reviewers for their helpful comments on the manuscript.

Transparency document. Supplementary material

Transparency document associated with this article can be found in the online version at <https://doi.org/10.1016/j.dib.2018.10.164>.

Appendix A. Supplementary material

Supplementary data associated with this article can be found in the online version at <https://doi.org/10.1016/j.dib.2018.10.164>.

References

- [1] H. Ehrlich, K.D. Demadis, O.S. Pokrovsky, P.G. Koutsoukos, Modern views on desilicification: biosilica and abiotic silica dissolution in natural and artificial environments, *Chem. Rev.* 110 (2010) 4656–4689.
- [2] M. Sommer, D. Kaczorek, Y. Kuzyakov, J. Breuer, Silicon pools and fluxes in soils and landscapes – a review, *J. Plant Nutr. Soil Sci.* 169 (2006) 310–329.
- [3] D. Puppe, O. Ehrmann, D. Kaczorek, M. Wanner, M. Sommer, The protozoic Si pool in temperate forest ecosystems – Quantification, abiotic controls and interactions with earthworms, *Geoderma* 243–244 (2015) 196–204.
- [4] D.J. Conley, Terrestrial ecosystems and the global biogeochemical silica cycle, *Glob. Biogeochem. Cycles* 16 (2002) 68.1–68.8.
- [5] F.A. Street-Perrott, P.A. Barker, Biogenic silica: a neglected component of the coupled global continental biogeochemical cycles of carbon and silicon, *Earth Surf. Process. Landforms* 33 (2008) 1436–1457.
- [6] E. Struyf, A. Smis, S. Van Damme, P. Meire, D.J. Conley, The global biogeochemical silicon cycle, *Silicon* 1 (2009) 207–213.
- [7] F. Bartoli, The biogeochemical cycle of silicon in two temperate forest ecosystems, *Environ. Biogeochem.: Ecol. Bull.* 35 (1983) 469–476.
- [8] J.D. Meunier, F. Colin, C. Alarcon, Biogenic silica storage in soils, *Geology* 27 (1999) 835–838.
- [9] J.C. Carey, R.W. Fulweiler, The terrestrial silica pump, *PLoS One* 7 (2012) e52932.
- [10] Y. Aoki, M. Hoshino, T. Matsubara, Silica and testate amoebae in a soil under pine–oak forest, *Geoderma* 142 (2007) 29–35.
- [11] M. Sommer, H. Jochheim, A. Höhn, J. Breuer, Z. Zagorski, J. Busse, D. Barkusky, K. Meier, D. Puppe, M. Wanner, D. Kaczorek, Si cycling in a forest biogeosystem – the importance of transient state biogenic Si pools, *Biogeosciences* 10 (2013) 4991–5007.
- [12] D. Puppe, D. Kaczorek, M. Wanner, M. Sommer, Dynamics and drivers of the protozoic Si pool along a 10-year chronosequence of initial ecosystem states, *Ecol. Eng.* 70 (2014) 477–482.
- [13] D. Puppe, A. Höhn, D. Kaczorek, M. Wanner, M. Sommer, As time goes by – spatiotemporal changes of biogenic Si pools in initial soils of an artificial catchment in NE Germany, *Appl. Soil Ecol.* 105 (2016) 9–16.
- [14] D. Puppe, A. Höhn, D. Kaczorek, M. Wanner, M. Wehrhan, M. Sommer, How big is the influence of biogenic silicon pools on short-term changes in water-soluble silicon in soils? Implications from a study of a 10-year-old soil-plant system, *Biogeosciences* 14 (2017) 5239–5252.
- [15] A.L. Creevy, J. Fisher, D. Puppe, D.M. Wilkinson, Protist diversity on a nature reserve in NW England – with particular reference to their role in soil biogenic silicon pools, *Pedobiologia* 59 (2016) 51–59.
- [16] R. Meisterfeld, Testate amoebae with filopodia, in: J.J. Lee, G.F. Leedale, P. Bradbury (Eds.), *The Illustrated Guide to the Protozoa*, Society of Protozoologists, Lawrence, KS, USA, 2002, pp. 1054–1084.
- [17] WRB, *World Reference Base for Soil Resources 2006 – World Soil Resources Report No. 103*. FAO, Rome, 2006.
- [18] W. Gerwin, W. Schaaf, D. Biemelt, A. Fischer, S. Winter, R.F. Hüttel, The artificial catchment “Chicken Creek” (Lusatia, Germany) – a landscape laboratory for interdisciplinary studies of initial ecosystem development, *Ecol. Eng.* 35 (2009) 1786–1796.
- [19] G. Kendzia, R. Reißmann, T. Neumann, Targeted development of wetland habitats for nature conservation fed by natural inflow in the post-mining landscape of Lusatia, *World Min.* 60 (2008) 88–95.
- [20] A. Höhn, M. Sommer, D. Kaczorek, G. Schalitz, J. Breuer, Silicon fractions in Histosols and Gleysols of a temperate grassland site, *J. Plant Nutr. Soil Sci.* 171 (2008) 409–418.
- [21] W. Foissner, Soil protozoa as bioindicators: pros and cons, methods, diversity, representative examples, *Agric. Ecosyst. Environ.* 74 (1999) 95–112.