




Universität Potsdam



Christian Bär | Nicolas Ginoux

Classical and Quantum Fields on Lorentzian Manifolds

Preprints des Instituts für Mathematik der Universität Potsdam
I (2012) 15

Christian Bär | Nicolas Ginoux

Classical and Quantum Fields on Lorentzian Manifolds

Bibliografische Information der Deutschen Nationalbibliothek

Die Deutsche Nationalbibliothek verzeichnet diese Publikation in der Deutschen Nationalbibliografie; detaillierte bibliografische Daten sind im Internet über <http://dnb.de> abrufbar.

Universitätsverlag Potsdam 2012

<http://info.ub.uni-potsdam.de/verlag.htm>

Am Neuen Palais 10, 14469 Potsdam
Tel.: +49 (0)331 977 2533 / Fax: 2292
E-Mail: verlag@uni-potsdam.de

Die Schriftenreihe **Preprints des Instituts für Mathematik der Universität Potsdam** wird herausgegeben vom Institut für Mathematik der Universität Potsdam.

ISSN (online) 2193-6943

Kontakt:

Institut für Mathematik
Am Neuen Palais 10
14469 Potsdam
Tel.: +49 (0)331 977 1028
WWW: <http://www.math.uni-potsdam.de>

Titelabbildungen:

1. Karla Fritze | Institutsgebäude auf dem Campus Neues Palais
 2. Nicolas Curien, Wendelin Werner | Random hyperbolic triangulation
- Published at: <http://arxiv.org/abs/1105.5089>
Das Manuskript ist urheberrechtlich geschützt.

Online veröffentlicht auf dem Publikationsserver der Universität Potsdam

URL <http://pub.ub.uni-potsdam.de/volltexte/2012/5997/>

URN <urn:nbn:de:kobv:517-opus-59973>

<http://nbn-resolving.de/urn:nbn:de:kobv:517-opus-59973>

CLASSICAL AND QUANTUM FIELDS ON LORENTZIAN MANIFOLDS

CHRISTIAN BÄR AND NICOLAS GINOUX

ABSTRACT. We construct bosonic and fermionic locally covariant quantum field theories on curved backgrounds for large classes of fields. We investigate the quantum field and n -point functions induced by suitable states.

1. INTRODUCTION

Classical fields on spacetime are mathematically modeled by sections of a vector bundle over a Lorentzian manifold. The field equations are usually partial differential equations. We introduce a class of differential operators, called Green-hyperbolic operators, which have good analytical solubility properties. This class includes wave operators as well as Dirac type operators.

In order to quantize such a classical field theory on a curved background, we need local algebras of observables. They come in two flavors, bosonic algebras encoding the canonical commutation relations and fermionic algebras encoding the canonical anti-commutation relations. We show how such algebras can be associated to manifolds equipped with suitable Green-hyperbolic operators. We prove that we obtain locally covariant quantum field theories in the sense of [11]. There is a large literature where such constructions are carried out for particular examples of fields, see e.g. [14, 17, 18, 20, 26, 38]. In all these papers the well-posedness of the Cauchy problem plays an important role. We avoid using the Cauchy problem altogether and only make use of Green's operators. In this respect, our approach is similar to the one in [39]. This allows us to deal with larger classes of fields, see Section 2.7, and to treat them systematically. Much of the earlier work on constructing observable algebras for particular examples can be subsumed under this general approach.

It turns out that bosonic algebras can be obtained in much more general situations than fermionic algebras. For instance, for the classical Dirac field both constructions are possible. Hence, on the level of observable algebras, there is no spin-statistics theorem. In order to obtain results like Theorem 5.1 in [41] one needs more structure, namely representations of the observable algebras with good properties.

In order to produce numbers out of our quantum field theory that can be compared to experiments, we need states, in addition to observables. We show how states with suitable regularity properties give rise to quantum fields and n -point functions.

Date: April 29, 2011.

2010 Mathematics Subject Classification. 58J45,35Lxx,81T20.

Key words and phrases. Wave operator, Dirac-type operator, globally hyperbolic spacetime, Green's operator, CCR-algebra, CAR-algebra, state, representation, locally covariant quantum field theory, quantum field, n -point function.

We check that they have the properties expected from traditional quantum field theories on a Minkowski background.

Acknowledgments. It is a pleasure to thank Alexander Strohmaier and Rainer Verch for very valuable discussion. The authors would also like to thank SPP 1154 “Globale Differentialgeometrie” and SFB 647 “Raum-Zeit-Materie”, both funded by Deutsche Forschungsgemeinschaft, for financial support.

2. FIELD EQUATIONS ON LORENTZIAN MANIFOLDS

2.1. Globally hyperbolic manifolds. We begin by fixing notation and recalling general facts about Lorentzian manifolds, see e.g. [30] or [4] for more details. Unless mentioned otherwise, the pair (M, g) will stand for a smooth m -dimensional manifold M equipped with a smooth Lorentzian metric g , where our convention for Lorentzian signature is $(- + \cdots +)$. The associated volume element will be denoted by dV . We shall also assume our Lorentzian manifold (M, g) to be time-orientable, i.e., that there exists a smooth timelike vector field on M . Time-oriented Lorentzian manifolds will be also referred to as *spacetimes*. Note that in contrast to conventions found elsewhere, we do not assume that a spacetime is connected nor do we assume that its dimension be $m = 4$.

For every subset A of a spacetime M we denote the causal future and past of A in M by $J_+(A)$ and $J_-(A)$, respectively. If we want to emphasize the ambient space M in which the causal future or past of A is considered, we write $J_{\pm}^M(A)$ instead of $J_{\pm}(A)$. Causal curves will always be implicitly assumed (future or past) oriented.

Definition 2.1. A *Cauchy hypersurface* in a spacetime (M, g) is a subset of M which is met exactly once by every inextendible timelike curve.

Cauchy hypersurfaces are always topological hypersurfaces but need not be smooth. All Cauchy hypersurfaces of a spacetime are homeomorphic.

Definition 2.2. A spacetime (M, g) is called *globally hyperbolic* if and only if it contains a Cauchy hypersurface.

A classical result of R. Geroch [21] says that a globally hyperbolic spacetime can be foliated by Cauchy hypersurfaces. It is a rather recent and very important result that this also holds in the smooth category:

Theorem 2.3 (A. Bernal and M. Sánchez [6, Thm. 1.1]). *Let (M, g) be a globally hyperbolic spacetime.*

Then there exists a smooth manifold Σ , a smooth one-parameter-family of Riemannian metrics $(g_t)_t$ on Σ and a smooth positive function β on $\mathbb{R} \times \Sigma$ such that (M, g) is isometric to $(\mathbb{R} \times \Sigma, -\beta dt^2 \oplus g_t)$. Each $\{t\} \times \Sigma$ corresponds to a smooth spacelike Cauchy hypersurface in (M, g) .

For our purposes, we shall need a slightly stronger version of Theorem 2.3 where one of the Cauchy hypersurfaces $\{t\} \times \Sigma$ can be prescribed:

Theorem 2.4 (A. Bernal and M. Sánchez [7, Thm. 1.2]). *Let (M, g) be a globally hyperbolic spacetime and $\tilde{\Sigma}$ a smooth spacelike Cauchy hypersurface in (M, g) . Then there exists a smooth splitting $(M, g) \cong (\mathbb{R} \times \Sigma, -\beta dt^2 \oplus g_t)$ as in Theorem 2.3 such that $\tilde{\Sigma}$ corresponds to $\{0\} \times \Sigma$.*

We shall also need the following result which tells us that one can extend any compact acausal spacelike submanifold to a smooth spacelike Cauchy hypersurface.

Here a subset of a spacetime is called *acausal* if no causal curve meets it more than once.

Theorem 2.5 (A. Bernal and M. Sánchez [7, Thm. 1.1]). *Let (M, g) be a globally hyperbolic spacetime and let $K \subset M$ be a compact acausal smooth spacelike submanifold with boundary.*

Then there exists a smooth spacelike Cauchy hypersurface Σ in (M, g) with $K \subset \Sigma$.

Definition 2.6. A closed subset $A \subset M$ is called *spacelike compact* if there exists a compact subset $K \subset M$ such that $A \subset J^M(K) := J_-^M(K) \cup J_+^M(K)$.

Note that a spacelike compact subset is in general not compact, but its intersection with any Cauchy hypersurface is compact, see e.g. [4, Cor. A.5.4].

Definition 2.7. A subset Ω of a spacetime M is called *causally compatible* if and only if $J_\pm^\Omega(x) = J_\pm^M(x) \cap \Omega$ for every $x \in \Omega$.

This means that every causal curve joining two points in Ω must be contained entirely in Ω .

2.2. Differential operators and Green's functions. A *differential operator* of order (at most) k on a vector bundle $S \rightarrow M$ over $\mathbb{K} = \mathbb{R}$ or $\mathbb{K} = \mathbb{C}$ is a linear map $P : C^\infty(M, S) \rightarrow C^\infty(M, S)$ which in local coordinates $x = (x^1, \dots, x^m)$ of M and with respect to a local trivialization looks like

$$P = \sum_{|\alpha| \leq k} A_\alpha(x) \frac{\partial^\alpha}{\partial x^\alpha}.$$

Here $C^\infty(M, S)$ denotes the space of smooth sections of $S \rightarrow M$, $\alpha = (\alpha_1, \dots, \alpha_m) \in \mathbb{N}_0 \times \dots \times \mathbb{N}_0$ runs over multi-indices, $|\alpha| = \alpha_1 + \dots + \alpha_m$ and $\frac{\partial^\alpha}{\partial x^\alpha} = \frac{\partial^{|\alpha|}}{\partial (x^1)^{\alpha_1} \dots \partial (x^m)^{\alpha_m}}$. The *principal symbol* σ_P of P associates to each covector $\xi \in T_x^*M$ a linear map $\sigma_P(\xi) : S_x \rightarrow S_x$. Locally, it is given by

$$\sigma_P(\xi) = \sum_{|\alpha|=k} A_\alpha(x) \xi^\alpha$$

where $\xi^\alpha = \xi_1^{\alpha_1} \dots \xi_m^{\alpha_m}$ and $\xi = \sum_j \xi_j dx^j$. If P and Q are two differential operators of order k and ℓ respectively, then $Q \circ P$ is a differential operator of order $k + \ell$ and

$$\sigma_{Q \circ P}(\xi) = \sigma_Q(\xi) \circ \sigma_P(\xi).$$

For any linear differential operator $P : C^\infty(M, S) \rightarrow C^\infty(M, S)$ there is a unique formally dual operator $P^* : C^\infty(M, S^*) \rightarrow C^\infty(M, S^*)$ of the same order characterized by

$$\int_M \langle \varphi, P\psi \rangle dV = \int_M \langle P^*\varphi, \psi \rangle dV$$

for all $\psi \in C^\infty(M, S)$ and $\varphi \in C^\infty(M, S^*)$ with $\text{supp}(\varphi) \cap \text{supp}(\psi)$ compact. Here $\langle \cdot, \cdot \rangle : S^* \otimes S \rightarrow \mathbb{K}$ denotes the canonical pairing, i.e., the evaluation of a linear form in S_x^* on an element of S_x , where $x \in M$. We have $\sigma_{P^*}(\xi) = (-1)^k \sigma_P(\xi)^*$ where k is the order of P .

Definition 2.8. Let a vector bundle $S \rightarrow M$ be endowed with a non-degenerate inner product $\langle \cdot, \cdot \rangle$. A linear differential operator P on S is called *formally self-adjoint* if and only if

$$\int_M \langle P\varphi, \psi \rangle dV = \int_M \langle \varphi, P\psi \rangle dV$$

holds for all $\varphi, \psi \in C^\infty(M, S)$ with $\text{supp}(\varphi) \cap \text{supp}(\psi)$ compact. Similarly, we call P *formally skew-adjoint* if instead

$$\int_M \langle P\varphi, \psi \rangle dV = - \int_M \langle \varphi, P\psi \rangle dV.$$

We recall the definition of advanced and retarded Green's operators for a linear differential operator.

Definition 2.9. Let P be a linear differential operator acting on the sections of a vector bundle S over a Lorentzian manifold M . An *advanced Green's operator* for P on M is a linear map

$$G_+ : C_c^\infty(M, S) \rightarrow C^\infty(M, S)$$

satisfying:

- (G₁) $P \circ G_+ = \text{id}_{C_c^\infty(M, S)}$;
- (G₂) $G_+ \circ P|_{C_c^\infty(M, S)} = \text{id}_{C_c^\infty(M, S)}$;
- (G₃⁺) $\text{supp}(G_+\varphi) \subset J_+^M(\text{supp}(\varphi))$ for any $\varphi \in C_c^\infty(M, S)$.

A *retarded Green's operator* for P on M is a linear map $G_- : C_c^\infty(M, S) \rightarrow C^\infty(M, S)$ satisfying (G₁), (G₂), and

- (G₃⁻) $\text{supp}(G_-\varphi) \subset J_-^M(\text{supp}(\varphi))$ for any $\varphi \in C_c^\infty(M, S)$.

Here we denote by $C_c^\infty(M, S)$ the space of compactly supported smooth sections of S .

Definition 2.10. Let $P : C^\infty(M, S) \rightarrow C^\infty(M, S)$ be a linear differential operator. We call P *Green-hyperbolic* if the restriction of P to any globally hyperbolic subregion of M has advanced and retarded Green's operators.

Remark 2.11. If the Green's operators of the restriction of P to a globally hyperbolic subregion exist, then they are necessarily unique, see Remark 3.7.

2.3. Wave operators. The most prominent class of Green-hyperbolic operators are wave operators, sometimes also called normally hyperbolic operators.

Definition 2.12. A linear differential operator of second order $P : C^\infty(M, S) \rightarrow C^\infty(M, S)$ is called a *wave operator* if its principal symbol is given by the Lorentzian metric, i.e., for all $\xi \in T^*M$ we have

$$\sigma_P(\xi) = -\langle \xi, \xi \rangle \cdot \text{id}.$$

In other words, if we choose local coordinates x^1, \dots, x^m on M and a local trivialization of S , then

$$P = - \sum_{i,j=1}^m g^{ij}(x) \frac{\partial^2}{\partial x^i \partial x^j} + \sum_{j=1}^m A_j(x) \frac{\partial}{\partial x^j} + B(x)$$

where A_j and B are matrix-valued coefficients depending smoothly on x and (g^{ij}) is the inverse matrix of (g_{ij}) with $g_{ij} = \langle \frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j} \rangle$. If P is a wave operator, then so is its dual operator P^* . In [4, Cor. 3.4.3] it has been shown that wave operators are Green-hyperbolic.

Example 2.13 (d'Alembert operator). Let S be the trivial line bundle so that sections of S are just functions. The d'Alembert operator $P = \square = -\text{div} \circ \text{grad}$ is a formally self-adjoint wave operator, see e.g. [4, p. 26].

Example 2.14 (connection-d'Alembert operator). More generally, let S be a vector bundle and let ∇ be a connection on S . This connection and the Levi-Civita connection on T^*M induce a connection on $T^*M \otimes S$, again denoted ∇ . We define the connection-d'Alembert operator \square^∇ to be the composition of the following three maps

$$C^\infty(M, S) \xrightarrow{\nabla} C^\infty(M, T^*M \otimes S) \xrightarrow{\nabla} C^\infty(M, T^*M \otimes T^*M \otimes S) \xrightarrow{-\text{tr} \otimes \text{id}_S} C^\infty(M, S)$$

where $\text{tr} : T^*M \otimes T^*M \rightarrow \mathbb{R}$ denotes the metric trace, $\text{tr}(\xi \otimes \eta) = \langle \xi, \eta \rangle$. We compute the principal symbol,

$$\sigma_{\square^\nabla}(\xi)\varphi = -(\text{tr} \otimes \text{id}_S) \circ \sigma_\nabla(\xi) \circ \sigma_\nabla(\xi)(\varphi) = -(\text{tr} \otimes \text{id}_S)(\xi \otimes \xi \otimes \varphi) = -\langle \xi, \xi \rangle \varphi.$$

Hence \square^∇ is a wave operator.

Example 2.15 (Hodge-d'Alembert operator). Let $S = \Lambda^k T^*M$ be the bundle of k -forms. Exterior differentiation $d : C^\infty(M, \Lambda^k T^*M) \rightarrow C^\infty(M, \Lambda^{k+1} T^*M)$ increases the degree by one while the codifferential $\delta = d^* : C^\infty(M, \Lambda^k T^*M) \rightarrow C^\infty(M, \Lambda^{k-1} T^*M)$ decreases the degree by one. While d is independent of the metric, the codifferential δ does depend on the Lorentzian metric. The operator $P = -d\delta - \delta d$ is a formally self-adjoint wave operator.

2.4. The Proca equation. The Proca operator is an example of a Green-hyperbolic operator of second order which is not a wave operator. First we need the following observation:

Lemma 2.16. *Let M be globally hyperbolic, let $S \rightarrow M$ be a vector bundle and let P and Q be differential operators acting on sections of S . Suppose P has advanced and retarded Green's operators G_+ and G_- .*

If Q commutes with P , then it also commutes with G_+ and with G_- .

Proof. Assume $[P, Q] = 0$. We consider

$$\tilde{G}_\pm := G_\pm + [G_\pm, Q] : C_c^\infty(M, s) \rightarrow C_{\text{sc}}^\infty(M, S).$$

We compute on $C_c^\infty(M, S)$:

$$\tilde{G}_\pm P = G_\pm P + G_\pm QP - QG_\pm P = \text{id} + G_\pm PQ - Q = \text{id} + Q - Q = \text{id}$$

and similarly $P\tilde{G}_\pm = \text{id}$. Hence \tilde{G}_\pm are also advanced and retarded Green's operators, respectively. By Remark 2.11, Green's operators are unique, hence $\tilde{G}_\pm = G_\pm$ and therefore $[G_\pm, Q] = 0$. \square

Example 2.17 (Proca operator). The discussion of this example follows [39, p. 116f], see also [20] where the discussion is based on the Cauchy problem. The Proca equation describes massive vector bosons. We take $S = T^*M$ and let $m_0 > 0$. The Proca equation is

$$(1) \quad P\varphi := \delta d\varphi + m_0^2\varphi = 0$$

where $\varphi \in C^\infty(M, S)$. Applying δ to (1) we obtain, using $\delta^2 = 0$ and $m_0 \neq 0$,

$$(2) \quad \delta\varphi = 0$$

and hence

$$(3) \quad (d\delta + \delta d)\varphi + m_0^2\varphi = 0.$$

Conversely, (2) and (3) clearly imply (1).

Since $\tilde{P} := d\delta + \delta d + m_0^2$ is minus a wave operator, it has Green's operators \tilde{G}_\pm . We define

$$G_\pm : C_c^\infty(M, S) \rightarrow C_{\text{sc}}^\infty(M, S), \quad G_\pm := (m_0^{-2}d\delta + \text{id}) \circ \tilde{G}_\pm = \tilde{G}_\pm \circ (m_0^{-2}d\delta + \text{id}).$$

The last equality holds because d and δ commute with \tilde{P} . For $\varphi \in C_c^\infty(M, S)$ we compute

$$G_\pm P\varphi = \tilde{G}_\pm(m_0^{-2}d\delta + \text{id})(\delta d + m_0^2)\varphi = \tilde{G}_\pm \tilde{P}\varphi = \varphi$$

and similarly $PG_\pm\varphi = \varphi$. Since the differential operator $m_0^{-2}d\delta + \text{id}$ does not increase supports, the third axiom in the definition of advanced and retarded Green's operators holds as well.

This shows that G_+ and G_- are advanced and retarded Green's operators for P , respectively. Thus P is not a wave operator but Green-hyperbolic.

2.5. Dirac type operators. The most important Green-hyperbolic operators of first order are the so-called Dirac type operators.

Definition 2.18. A linear differential operator $D : C^\infty(M, S) \rightarrow C^\infty(M, S)$ of first order is called *of Dirac type*, if $-D^2$ is a wave operator.

Remark 2.19. If D is of Dirac type, then i times its principal symbol satisfies the Clifford relations

$$(i\sigma_D(\xi))^2 = -\sigma_{D^2}(\xi) = -\langle \xi, \xi \rangle \cdot \text{id},$$

hence by polarization

$$(i\sigma_D(\xi))(i\sigma_D(\eta)) + (i\sigma_D(\eta))(i\sigma_D(\xi)) = -2\langle \xi, \eta \rangle \cdot \text{id}.$$

The bundle S thus becomes a module over the bundle of Clifford algebras $\text{Cl}(TM)$ associated with $(TM, \langle \cdot, \cdot \rangle)$. See [5, Sec. 1.1] or [27, Ch. I] for the definition and properties of the Clifford algebra $\text{Cl}(V)$ associated with a vector space V with inner product.

Remark 2.20. If D is of Dirac type, then so is its dual operator D^* . On a globally hyperbolic region let G_+ be the advanced Green's operator for D^2 which exists since $-D^2$ is a wave operator. Then it is not hard to check that $D \circ G_+$ is an advanced Green's operator for D , see e.g. the proof of Theorem 2.3 in [14] or [29, Thm. 3.2]. The same discussion applies to the retarded Green's operator. Hence any Dirac type operator is Green-hyperbolic.

Example 2.21 (Classical Dirac operator). If the spacetime M carries a spin structure, then one can define the spinor bundle $S = \Sigma M$ and the classical Dirac operator

$$D : C^\infty(M, \Sigma M) \rightarrow C^\infty(M, \Sigma M), \quad D\varphi := i \sum_{j=1}^m \varepsilon_j e_j \cdot \nabla_{e_j} \varphi.$$

Here $(e_j)_{1 \leq j \leq m}$ is a local orthonormal basis of the tangent bundle, $\varepsilon_j = \langle e_j, e_j \rangle = \pm 1$ and “ \cdot ” denotes the Clifford multiplication, see e.g. [5] or [3, Sec. 2]. The principal symbol of D is given by

$$\sigma_D(\xi)\psi = i\xi^\sharp \cdot \psi.$$

Here ξ^\sharp denotes the tangent vector dual to the 1-form ξ via the Lorentzian metric, i.e., $\langle \xi^\sharp, Y \rangle = \xi(Y)$ for all tangent vectors Y over the same point of the manifold. Hence

$$\sigma_{D^2}(\xi)\psi = \sigma_D(\xi)\sigma_D(\xi)\psi = -\xi^\sharp \cdot \xi^\sharp \cdot \psi = \langle \xi, \xi \rangle \psi.$$

Thus $P = -D^2$ is a wave operator. Moreover, D is formally self-adjoint, see e.g. [3, p. 552].

Example 2.22 (Twisted Dirac operators). More generally, let $E \rightarrow M$ be a complex vector bundle equipped with a non-degenerate Hermitian inner product and a metric connection ∇^E over a spin spacetime M . In the notation of Example 2.21, one may define the Dirac operator of M twisted with E by

$$D^E := i \sum_{j=1}^m \varepsilon_j e_j \cdot \nabla_{e_j}^{\Sigma M \otimes E} : C^\infty(M, \Sigma M \otimes E) \rightarrow C^\infty(M, \Sigma M \otimes E),$$

where $\nabla^{\Sigma M \otimes E}$ is the tensor product connection on $\Sigma M \otimes E$. Again, D^E is a formally self-adjoint Dirac type operator.

Example 2.23 (Euler operator). In Example 2.15, replacing $\Lambda^k T^*M$ by $S := \Lambda T^*M \otimes \mathbb{C} = \bigoplus_{k=0}^n \Lambda^k T^*M \otimes \mathbb{C}$, the Euler operator $D = i(d - \delta)$ defines a formally self-adjoint Dirac type operator. In case M is spin, the Euler operator coincides with the Dirac operator of M twisted with ΣM .

Example 2.24 (Buchdahl operators). On a 4-dimensional spin spacetime M , consider the standard orthogonal and parallel splitting $\Sigma M = \Sigma_+ M \oplus \Sigma_- M$ of the complex spinor bundle of M into spinors of positive and negative chirality. The finite dimensional irreducible representations of the simply-connected Lie group $\text{Spin}^0(3, 1)$ are given by $\Sigma_+^{(k/2)} \otimes \Sigma_-^{(\ell/2)}$ where $k, \ell \in \mathbb{N}$. Here $\Sigma_+^{(k/2)} = \Sigma_+^{\odot k}$ is the k -th symmetric tensor product of the positive half-spinor representation Σ_+ and similarly for $\Sigma_-^{(\ell/2)}$. Let the associated vector bundles $\Sigma_\pm^{(k/2)} M$ carry the induced inner product and connection.

For $s \in \mathbb{N}$, $s \geq 1$, consider the twisted Dirac operator $D^{(s)}$ acting on sections of $\Sigma M \otimes \Sigma_+^{((s-1)/2)} M$. In the induced splitting

$$\Sigma M \otimes \Sigma_+^{((s-1)/2)} M = \Sigma_+ M \otimes \Sigma_+^{(s-1/2)} M \oplus \Sigma_- M \otimes \Sigma_+^{((s-1)/2)} M$$

the operator $D^{(s)}$ is of the form

$$\begin{pmatrix} 0 & D_-^{(s)} \\ D_+^{(s)} & 0 \end{pmatrix}$$

because Clifford multiplication by vectors exchanges the chiralities. The Clebsch-Gordan formulas [10, Prop. II.5.5] tell us that the representation $\Sigma_+ \otimes \Sigma_+^{((s-1)/2)}$ splits as

$$\Sigma_+ \otimes \Sigma_+^{((s-1)/2)} = \Sigma_+^{(s/2)} \oplus \Sigma_+^{(s/2-1)}.$$

Hence we have the corresponding parallel orthogonal projections

$$\pi_s : \Sigma_+ M \otimes \Sigma_+^{((s-1)/2)} M \rightarrow \Sigma_+^{(s/2)} M \quad \text{and} \quad \pi'_s : \Sigma_+ M \otimes \Sigma_+^{((s-1)/2)} M \rightarrow \Sigma_+^{(s/2-1)} M.$$

On the other hand, the representation $\Sigma_- \otimes \Sigma_+^{((s-1)/2)}$ is irreducible. Now *Buchdahl operators* are the operators of the form

$$B_{\mu_1, \mu_2, \mu_3}^{(s)} := \begin{pmatrix} \mu_1 \cdot \pi_s + \mu_2 \cdot \pi'_s & D_-^{(s)} \\ D_+^{(s)} & \mu_3 \cdot \text{id} \end{pmatrix}$$

where $\mu_1, \mu_2, \mu_3 \in \mathbb{C}$ are constants. By definition, $B_{\mu_1, \mu_2, \mu_3}^{(s)}$ is of the form $D^{(s)} + b$, where b is of order zero. In particular, $B_{\mu_1, \mu_2, \mu_3}^{(s)}$ is a Dirac-type operator, hence it is Green-hyperbolic.

If M were Riemannian, then $D^{(s)}$ would be formally self-adjoint. Hence the operator $B_{\mu_1, \mu_2, \mu_3}^{(s)}$ would be formally self-adjoint if and only if the constants μ_1, μ_2, μ_3 are real. In Lorentzian signature, $\Sigma_+ M$ and $\Sigma_- M$ are isotropic for the natural inner product on ΣM , so that the bundles on which the Buchdahl operators act, carry no natural non-degenerate inner product.

For a definition of Buchdahl operators using indices we refer to [12, 13, 44] and to [28, Def. 8.1.4, p. 104].

2.6. The Rarita-Schwinger operator. For the Rarita-Schwinger operator on Riemannian manifolds, we refer to [43, Sec. 2], see also [8, Sec. 2]. In this section let the spacetime M be spin and consider the Clifford-multiplication $\gamma : T^*M \otimes \Sigma M \rightarrow \Sigma M$, $\theta \otimes \psi \mapsto \theta^\sharp \cdot \psi$, where ΣM is the complex spinor bundle of M . Then there is the representation theoretic splitting of $T^*M \otimes \Sigma M$ into the orthogonal and parallel sum

$$T^*M \otimes \Sigma M = \iota(\Sigma M) \oplus \Sigma^{3/2}M,$$

where $\Sigma^{3/2}M := \ker(\gamma)$ and $\iota(\psi) := -\frac{1}{m} \sum_{j=1}^m e_j^* \otimes e_j \cdot \psi$. Here again $(e_j)_{1 \leq j \leq m}$ is a local orthonormal basis of the tangent bundle. Let \mathcal{D} be the twisted Dirac operator on $T^*M \otimes \Sigma M$, that is, $\mathcal{D} := i \cdot (\text{id} \otimes \gamma) \circ \nabla$, where ∇ denotes the induced covariant derivative on $T^*M \otimes \Sigma M$.

Definition 2.25. The Rarita-Schwinger operator on the spin spacetime M is defined by $\mathcal{Q} := (\text{id} - \iota \circ \gamma) \circ \mathcal{D} : C^\infty(M, \Sigma^{3/2}M) \rightarrow C^\infty(M, \Sigma^{3/2}M)$.

By definition, the Rarita-Schwinger operator is pointwise obtained as the orthogonal projection onto $\Sigma^{3/2}M$ of the twisted Dirac operator \mathcal{D} restricted to a section of $\Sigma^{3/2}M$. Using the above formula for ι , the Rarita-Schwinger operator can be written down explicitly:

$$\mathcal{Q}\psi = i \cdot \sum_{\beta=1}^m e_\beta^* \otimes \sum_{\alpha=1}^m \varepsilon_\alpha (e_\alpha \cdot \nabla_{e_\alpha} \psi_\beta - \frac{2}{m} e_\beta \cdot \nabla_{e_\alpha} \psi_\alpha)$$

for all $\psi = \sum_{\beta=1}^m e_\beta^* \otimes \psi_\beta \in C^\infty(M, \Sigma^{3/2}M)$, where here ∇ is the standard connection on ΣM . It can be checked that \mathcal{Q} is a formally self-adjoint linear differential operator of first order, with principal symbol

$$\sigma_{\mathcal{Q}}(\xi) : \psi \mapsto i \left\{ (\text{id} \otimes \xi^\sharp \cdot) \psi - \frac{2}{m} \sum_{\beta=1}^m e_\beta^* \otimes e_\beta \cdot (\xi^\sharp \lrcorner \psi) \right\},$$

for all $\psi = \sum_{\beta=1}^m e_\beta^* \otimes \psi_\beta \in \Sigma^{3/2}M$. Here $X \lrcorner \psi$ denotes the insertion of the tangent vector X in the first factor, that is, $X \lrcorner \psi := \sum_{\beta=1}^m e_\beta^*(X) \psi_\beta$.

Lemma 2.26. *Let M be a spin spacetime of dimension $m \geq 3$. Then the characteristic variety of the Rarita-Schwinger operator of M coincides with the set of lightlike covectors.*

Proof. By definition, the characteristic variety of \mathcal{Q} is the set of nonzero covectors ξ for which $\sigma_{\mathcal{Q}}(\xi)$ is not invertible. Fix an arbitrary point $x \in M$. Let $\xi \in T_x^*M \setminus \{0\}$ be non-lightlike. Without loss of generality we may assume

that ξ is normalized and that the Lorentz orthonormal basis is chosen so that $\xi^\sharp = e_1$. Hence $\varepsilon_1 = 1$ if ξ is spacelike and $\varepsilon_1 = -1$ if ξ is timelike. Take $\psi = \sum_{\beta=1}^m e_\beta^* \otimes \psi_\beta \in \ker(\sigma_{\mathcal{Q}}(\xi))$. Then

$$\begin{aligned} 0 &= \sum_{\beta=1}^m e_\beta^* \otimes e_1 \cdot \psi_\beta - \frac{2}{m} \sum_{\beta=1}^m e_\beta^* \otimes e_\beta \cdot \psi_1 \\ &= \sum_{\beta=1}^m e_\beta^* \otimes \left(e_1 \cdot \psi_\beta - \frac{2}{m} e_\beta \cdot \psi_1 \right), \end{aligned}$$

which implies $e_1 \cdot \psi_\beta = \frac{2}{m} e_\beta \cdot \psi_1$ for all $\beta \in \{1, \dots, m\}$. Choosing $\beta = 1$, we obtain $e_1 \cdot \psi_1 = 0$ because $m \geq 3$. Hence $\psi_1 = 0$, from which $\psi_\beta = 0$ follows for all $\beta \in \{1, \dots, m\}$. Hence $\psi = 0$ and $\sigma_{\mathcal{Q}}(\xi)$ is invertible.

If $\xi \in T_x^*M \setminus \{0\}$ is lightlike, then we may assume that $\xi^\sharp = e_1 + e_2$, where $\varepsilon_1 = -1$ and $\varepsilon_2 = 1$. Choose $\psi_1 \in \Sigma_x M \setminus \{0\}$ with $(e_1 + e_2) \cdot \psi_1 = 0$. Such a ψ_1 exists because Clifford multiplication by a lightlike vector is nilpotent. Set $\psi_2 := -\psi_1$ and $\psi := e_1^* \otimes \psi_1 + e_2^* \otimes \psi_2$. Then $\psi \in \Sigma_x^{3/2}M \setminus \{0\}$ and

$$-i\sigma_{\mathcal{Q}}(\xi)(\psi) = \sum_{j=1}^2 e_j^* \otimes \underbrace{(e_1 + e_2) \cdot \psi_j}_{=0} - \frac{2}{m} e_j^* \otimes e_j \cdot \underbrace{(\psi_1 + \psi_2)}_{=0} = 0.$$

This shows $\psi \in \ker(\sigma_{\mathcal{Q}}(\xi))$ and hence $\sigma_{\mathcal{Q}}(\xi)$ is not invertible. \square

The same proof shows that in the Riemannian case the Rarita-Schwinger operator is elliptic.

Remark 2.27. Since the characteristic variety of the Rarita-Schwinger operator is exactly that of the Dirac operator, Lemma 2.26 together with [24, Thms. 23.2.4 & 23.2.7] imply that the Cauchy problem for \mathcal{Q} is well-posed in case M is globally hyperbolic. This implies they \mathcal{Q} has advanced and retarded Green's operators. Hence \mathcal{Q} is not of Dirac type but it is Green-hyperbolic.

Remark 2.28. The equations originally considered by Rarita and Schwinger in [33] correspond to the twisted Dirac operator \mathcal{D} restricted to $\Sigma^{3/2}M$ but not projected back to $\Sigma^{3/2}M$. In other words, they considered the operator

$$\mathcal{D}|_{C^\infty(M, \Sigma^{3/2}M)} : C^\infty(M, \Sigma^{3/2}M) \rightarrow C^\infty(M, T^*M \otimes \Sigma M).$$

These equations are over-determined. Therefore it is not a surprise that non-trivial solutions restrict the geometry of the underlying manifold as observed by Gibbons [22] and that this operator has no Green's operators.

2.7. Combining given operators into a new one. Given two Green-hyperbolic operators we can form the direct sum and obtain a new operator in a trivial fashion. It turns out that this operator is again Green-hyperbolic. Note that the two operators need not have the same order.

Lemma 2.29. *Let $S_1, S_2 \rightarrow M$ be two vector bundles over the globally hyperbolic manifold M . Let P_1 and P_2 be two Green-hyperbolic operators acting on sections of S_1 and S_2 respectively. Then*

$$P_1 \oplus P_2 := \begin{pmatrix} P_1 & 0 \\ 0 & P_2 \end{pmatrix} : C^\infty(M, S_1 \oplus S_2) \rightarrow C^\infty(M, S_1 \oplus S_2)$$

is Green-hyperbolic.

Proof. If G_1 and G_2 are advanced Green's operators for P_1 and P_2 respectively, then clearly $\begin{pmatrix} G_1 & 0 \\ 0 & G_2 \end{pmatrix}$ is an advanced Green's operator for $P_1 \oplus P_2$. The retarded case is analogous. \square

It is interesting to note that P_1 and P_2 need not have the same order. Hence Green-hyperbolic operators need not be hyperbolic in the usual sense. Moreover, it is not obvious that Green-hyperbolic operators have a well-posed Cauchy problem. For instance, if P_1 is a wave operator and P_2 a Dirac-type operator, then along a Cauchy hypersurface one would have to prescribe the normal derivative for the S_1 -component but not for the S_2 -component.

3. ALGEBRAS OF OBSERVABLES

Our next aim is to quantize the classical fields governed by Green-hyperbolic differential operators. We construct local algebras of observables and we prove that we obtain locally covariant quantum field theories in the sense of [11].

3.1. Bosonic quantization. In this section we show how a quantization process based on canonical commutation relations (CCR) can be carried out for formally self-adjoint Green-hyperbolic operators. This is a functorial procedure. We define the first category involved in the quantization process.

Definition 3.1. The category GlobHypGreen consists of the following objects and morphisms:

- An object in GlobHypGreen is a triple (M, S, P) , where
 - ▶ M is a globally hyperbolic spacetime,
 - ▶ S is a real vector bundle over M endowed with a non-degenerate inner product $\langle \cdot, \cdot \rangle$ and
 - ▶ P is a formally self-adjoint Green-hyperbolic operator acting on sections of S .
- A morphism between two objects (M_1, S_1, P_1) and (M_2, S_2, P_2) of GlobHypGreen is a pair (f, F) , where
 - ▶ f is a time-orientation preserving isometric embedding $M_1 \rightarrow M_2$ with $f(M_1)$ causally compatible and open in M_2 ,
 - ▶ F is a fiberwise isometric vector bundle isomorphism over f such that the following diagram commutes:

$$(4) \quad \begin{array}{ccc} C^\infty(M_2, S_2) & \xrightarrow{P_2} & C^\infty(M_2, S_2) \\ \text{res} \downarrow & & \downarrow \text{res} \\ C^\infty(M_1, S_1) & \xrightarrow{P_1} & C^\infty(M_1, S_1), \end{array}$$

where $\text{res}(\varphi) := F^{-1} \circ \varphi \circ f$ for every $\varphi \in C^\infty(M_2, S_2)$.

Note that morphisms exist only if the manifolds have equal dimension and the vector bundles have the same rank. Note furthermore, that the inner product $\langle \cdot, \cdot \rangle$ on S is not required to be positive or negative definite.

The causal compatibility condition, which is not automatically satisfied (see e.g. [4, Fig. 33]), ensures the commutation of the extension and restriction maps with the Green's operators:

Lemma 3.2. *Let (f, F) be a morphism between two objects (M_1, S_1, P_1) and (M_2, S_2, P_2) in the category GlobHypGreen and let $(G_1)_\pm$ and $(G_2)_\pm$ be the respective Green's operators for P_1 and P_2 . Denote by $\text{ext}(\varphi) \in C_c^\infty(M_2, S_2)$ the extension by 0 of $F \circ \varphi \circ f^{-1} : f(M_1) \rightarrow S_2$ to M_2 , for every $\varphi \in C_c^\infty(M_1, S_1)$. Then*

$$\text{res} \circ (G_2)_\pm \circ \text{ext} = (G_1)_\pm.$$

Proof. Set $(\widetilde{G}_1)_\pm := \text{res} \circ (G_2)_\pm \circ \text{ext}$ and fix $\varphi \in C_c^\infty(M_1, S_1)$. First observe that the causal compatibility condition on f implies that

$$\begin{aligned} \text{supp}((\widetilde{G}_1)_\pm(\varphi)) &= f^{-1}(\text{supp}((G_2)_\pm \circ \text{ext}(\varphi))) \\ &\subset f^{-1}(J_\pm^{M_2}(\text{supp}(\text{ext}(\varphi)))) \\ &= f^{-1}(J_\pm^{M_2}(f(\text{supp}(\varphi)))) \\ &= J_\pm^{M_1}(\text{supp}(\varphi)). \end{aligned}$$

In particular, $(\widetilde{G}_1)_\pm(\varphi)$ has spacelike compact support in M_1 and $(\widetilde{G}_1)_\pm$ satisfies Axiom (G_3) . Moreover, it follows from (4) that $P_2 \circ \text{ext} = \text{ext} \circ P_1$ on $C_c^\infty(M_1, S_1)$, which directly implies that $(\widetilde{G}_1)_\pm$ satisfies Axioms (G_1) and (G_2) as well. The uniqueness of the advanced and retarded Green's operators on M_1 yields $(\widetilde{G}_1)_\pm = (G_1)_\pm$. \square

Next we show how the Green's operators for a formally self-adjoint Green-hyperbolic operator provide a symplectic vector space in a canonical way. First we see how the Green's operators of an operator and of its formally dual operator are related.

Lemma 3.3. *Let M be a globally hyperbolic spacetime and G_+, G_- the advanced and retarded Green's operators for a Green-hyperbolic operator P acting on sections of $S \rightarrow M$. Then the advanced and retarded Green's operators G_+^* and G_-^* for P^* satisfy*

$$\int_M \langle G_\pm^* \varphi, \psi \rangle dV = \int_M \langle \varphi, G_\mp \psi \rangle dV$$

for all $\varphi \in C_c^\infty(M, S^*)$ and $\psi \in C_c^\infty(M, S)$.

Proof. Axiom (G_1) for the Green's operators implies that

$$\begin{aligned} \int_M \langle G_\pm^* \varphi, \psi \rangle dV &= \int_M \langle G_\pm^* \varphi, P(G_\mp \psi) \rangle dV \\ &= \int_M \langle P^*(G_\pm^* \varphi), G_\mp \psi \rangle dV \\ &= \int_M \langle \varphi, G_\mp \psi \rangle dV, \end{aligned}$$

where the integration by parts is justified since $\text{supp}(G_\pm^* \varphi) \cap \text{supp}(G_\mp \psi) \subset J_\pm^M(\text{supp}(\varphi)) \cap J_\mp^M(\text{supp}(\psi))$ is compact. \square

Proposition 3.4. *Let (M, S, P) be an object in the category GlobHypGreen . Set $G := G_+ - G_-$, where G_+, G_- are the advanced and retarded Green's operator for P , respectively.*

Then the pair $(\text{SYMPL}(M, S, P), \omega)$ is a symplectic vector space, where

$$\text{SYMPL}(M, S, P) := C_c^\infty(M, S) / \ker(G) \quad \text{and} \quad \omega([\varphi], [\psi]) := \int_M \langle G\varphi, \psi \rangle dV.$$

Here the square brackets $[\cdot]$ denote residue classes modulo $\ker(G)$.

Proof. The bilinear form $(\varphi, \psi) \mapsto \int_M \langle G\varphi, \psi \rangle dV$ on $C_c^\infty(M, S)$ is skew-symmetric as a consequence of Lemma 3.3 because P is formally self-adjoint. Its null-space is exactly $\ker(G)$. Therefore the induced bilinear form ω on the quotient space $\text{SYMPL}(M, S, P)$ is non-degenerate and hence a symplectic form. \square

Put $C_{\text{sc}}^\infty(M, S) := \{\varphi \in C_c^\infty(M, S) \mid \text{supp}(\varphi) \text{ is spacelike compact}\}$. The next result will in particular show that we can consider $\text{SYMPL}(M, S, P)$ as the space of smooth solutions of the equation $P\varphi = 0$ which have spacelike compact support.

Theorem 3.5. *Let M be a Lorentzian manifold, let $S \rightarrow M$ be a vector bundle, and let P be a Green-hyperbolic operator acting on sections of S . Let G_\pm be advanced and retarded Green's operators for P , respectively. Put*

$$G := G_+ - G_- : C_c^\infty(M, S) \rightarrow C_{\text{sc}}^\infty(M, S).$$

Then the following linear maps form a complex:

$$(5) \quad \{0\} \rightarrow C_c^\infty(M, S) \xrightarrow{P} C_c^\infty(M, S) \xrightarrow{G} C_{\text{sc}}^\infty(M, S) \xrightarrow{P} C_{\text{sc}}^\infty(M, S).$$

This complex is always exact at the first $C_c^\infty(M, S)$. If M is globally hyperbolic, then the complex is exact everywhere.

Proof. The proof follows the lines of [4, Thm. 3.4.7] where the result was shown for wave operators. First note that, by (G_3^\pm) in the definition of Green's operators, we have that $G_\pm : C_c^\infty(M, S) \rightarrow C_{\text{sc}}^\infty(M, S)$. It is clear from (G_1) and (G_2) that $PG = GP = 0$ on $C_c^\infty(M, S)$, hence (5) is a complex.

If $\varphi \in C_c^\infty(M, S)$ satisfies $P\varphi = 0$, then by (G_2) we have $\varphi = G_+P\varphi = 0$ which shows that $P|_{C_c^\infty(M, S)}$ is injective. Thus the complex is exact at the first $C_c^\infty(M, S)$.

From now on let M be globally hyperbolic. Let $\varphi \in C_c^\infty(M, S)$ with $G\varphi = 0$, i.e., $G_+\varphi = G_-\varphi$. We put $\psi := G_+\varphi = G_-\varphi \in C_c^\infty(M, S)$ and we see that $\text{supp}(\psi) = \text{supp}(G_+\varphi) \cap \text{supp}(G_-\varphi) \subset J_+(\text{supp}(\varphi)) \cap J_-(\text{supp}(\varphi))$. Since (M, g) is globally hyperbolic $J_+(\text{supp}(\varphi)) \cap J_-(\text{supp}(\varphi))$ is compact, hence $\psi \in C_c^\infty(M, S)$. From $P\psi = PG_+\varphi = \varphi$ we see that $\varphi \in P(C_c^\infty(M, S))$. This shows exactness at the second $C_c^\infty(M, S)$.

It remains to show that any $\varphi \in C_{\text{sc}}^\infty(M, S)$ with $P\varphi = 0$ is of the form $\varphi = G\psi$ with $\psi \in C_c^\infty(M, S)$. Using a cut-off function decompose φ as $\varphi = \varphi_+ - \varphi_-$ where $\text{supp}(\varphi_\pm) \subset J_\pm(K)$ where K is a suitable compact subset of M . Then $\psi := P\varphi_+ = P\varphi_-$ satisfies $\text{supp}(\psi) \subset J_+(K) \cap J_-(K)$. Thus $\psi \in C_c^\infty(M, S)$. We check that $G_+\psi = \varphi_+$. Namely, for all $\chi \in C_c^\infty(M, S^*)$ we have by Lemma 3.3

$$\int_M \langle \chi, G_+P\varphi_+ \rangle dV = \int_M \langle G_-^*\chi, P\varphi_+ \rangle dV = \int_M \langle P^*G_-^*\chi, \varphi_+ \rangle dV = \int_M \langle \chi, \varphi_+ \rangle dV.$$

The integration by parts in the second equality is justified because $\text{supp}(\varphi_+) \cap \text{supp}(G_-^*\chi) \subset J_+(K) \cap J_-(\text{supp}(\chi))$ is compact. Similarly, one shows $G_-\psi = \varphi_-$. Now $G\psi = G_+\psi - G_-\psi = \varphi_+ - \varphi_- = \varphi$ which concludes the proof. \square

In particular, given an object (M, S, P) in GlobHypGreen , the map G induces an isomorphism from

$$\text{SYMPL}(M, S, P) = C_c^\infty(M, S) / \ker(G) \xrightarrow{\cong} \ker(P) \cap C_{\text{sc}}^\infty(M, S).$$

Remark 3.6. Exactness at the first $C_c^\infty(M, S)$ in sequence (5) says that there are no non-trivial smooth solutions of $P\varphi = 0$ with compact support. Indeed, if M is globally hyperbolic, more is true.

If $\varphi \in C_c^\infty(M, S)$ solves $P\varphi = 0$ and $\text{supp}(\varphi)$ is future or past-compact, then $\varphi = 0$. Here a subset $A \subset M$ is called future-compact if $A \cap J_+(x)$ is compact for any $x \in M$. Past-compactness is defined similarly.

Proof. Let $\varphi \in C_c^\infty(M, S)$ solve $P\varphi = 0$ such that $\text{supp}(\varphi)$ is future-compact. For any $\chi \in C_c^\infty(M, S^*)$ we have

$$\int_M \langle \chi, \varphi \rangle dV = \int_M \langle P^* G_+^* \chi, \varphi \rangle dV = \int_M \langle G_+^* \chi, P\varphi \rangle dV = 0.$$

This shows $\varphi = 0$. The integration by parts is justified because $\text{supp}(G_+^* \chi) \cap \text{supp}(\varphi) \subset J_+(\text{supp}(\chi)) \cap \text{supp}(\varphi)$ is compact, see [4, Lemma A.5.3]. \square

Remark 3.7. Let M be a globally hyperbolic spacetime and (M, S, P) an object in GlobHypGreen . Then the Green's operators G_+ and G_- are unique. Namely, if G_+ and \tilde{G}_+ are advanced Green's operators for P , then for any $\varphi \in C_c^\infty(M, S)$ the section $\psi := G_+ \varphi - \tilde{G}_+ \varphi$ has past-compact support and satisfies $P\psi = 0$. By the previous remark, we have $\psi = 0$ which shows $G_+ = \tilde{G}_+$.

Now, let (f, F) be a morphism between two objects (M_1, S_1, P_1) and (M_2, S_2, P_2) in the category GlobHypGreen . For $\varphi \in C_c^\infty(M_1, S_1)$ consider the extension by zero $\text{ext}(\varphi) \in C_c^\infty(M_2, S_2)$ as in Lemma 3.2.

Lemma 3.8. Given a morphism (f, F) between two objects (M_1, S_1, P_1) and (M_2, S_2, P_2) in the category GlobHypGreen , extension by zero induces a symplectic linear map $\text{SYMPL}(f, F) : \text{SYMPL}(M_1, S_1, P_1) \rightarrow \text{SYMPL}(M_2, S_2, P_2)$.

Moreover,

$$(6) \quad \text{SYMPL}(\text{id}_M, \text{id}_S) = \text{id}_{\text{SYMPL}(M, S, P)}$$

and for any further morphism $(f', F') : (M_2, S_2, P_2) \rightarrow (M_3, S_3, P_3)$ one has

$$(7) \quad \text{SYMPL}((f', F') \circ (f, F)) = \text{SYMPL}(f', F') \circ \text{SYMPL}(f, F).$$

Proof. If $\varphi = P_1 \psi \in \ker(G_1) = P_1(C_c^\infty(M_1, S_1))$, then $\text{ext}(\varphi) = P_2(\text{ext}(\psi)) \in P_2(C_c^\infty(M_2, S_2)) = \ker(G_2)$. Hence ext induces a linear map

$$\text{SYMPL}(f, F) : C_c^\infty(M_1, S_1) / \ker(G_1) \rightarrow C_c^\infty(M_2, S_2) / \ker(G_2).$$

Furthermore, applying Lemma 3.2, we have, for any $\varphi, \psi \in C_c^\infty(M_1, S_1)$

$$\int_{M_2} \langle G_2(\text{ext}(\varphi)), \text{ext}(\psi) \rangle dV = \int_{M_1} \langle \text{res} \circ G_2 \circ \text{ext}(\varphi), \psi \rangle dV = \int_{M_1} \langle G_1 \varphi, \psi \rangle dV,$$

hence $\text{SYMPL}(f, F)$ is symplectic. Equation (6) is trivial and extending once or twice by 0 amounts to the same, so (7) holds as well. \square

Remark 3.9. Under the isomorphism $\text{SYMPL}(M, S, P) \rightarrow \ker(P) \cap C_{\text{sc}}^\infty(M, S)$ induced by G , the extension by zero corresponds to an extension as a smooth solution of $P\varphi = 0$ with spacelike compact support. This follows directly from Lemma 3.2.

In other words, for any morphism (f, F) from (M_1, S_1, P_1) to (M_2, S_2, P_2) in GlobHypGreen we have the following commutative diagram:

$$\begin{array}{ccc} \text{SYMPL}(M_1, S_1, P_1) & \xrightarrow{\text{SYMPL}(f, F)} & \text{SYMPL}(M_2, S_2, P_2) \\ \cong \downarrow & & \downarrow \cong \\ \ker(P_1) \cap C_{\text{sc}}^\infty(M_1, S_1) & \xrightarrow[\text{asolution}]{\text{extension as}} & \ker(P_2) \cap C_{\text{sc}}^\infty(M_2, S_2). \end{array}$$

Let Sympl denote the category of real symplectic vector spaces with symplectic linear maps as morphisms. Lemma 3.8 says that we have constructed a covariant functor

$$\text{SYMPL} : \text{GlobHypGreen} \longrightarrow \text{Sympl}.$$

In order to obtain an algebra-valued functor, we compose SYMPL with the functor CCR which associates to any symplectic vector space its Weyl algebra. Here “CCR” stands for “canonical commutation relations”. This is a general algebraic construction which is independent of the context of Green-hyperbolic operators and which is carried out in Section A.2. As a result, we obtain the functor

$$\mathfrak{A}_{\text{bos}} := \text{CCR} \circ \text{SYMPL} : \text{GlobHypGreen} \longrightarrow \text{C}^* \text{Alg},$$

where $\text{C}^* \text{Alg}$ is the category whose objects are the unital C^* -algebras and whose morphisms are the injective unit-preserving C^* -morphisms.

In the remainder of this section we show that the functor $\text{CCR} \circ \text{SYMPL}$ is a bosonic locally covariant quantum field theory. We call two subregions M_1 and M_2 of a spacetime M *causally disjoint* if and only if $J^M(M_1) \cap M_2 = \emptyset$. In other words, there are no causal curves joining M_1 and M_2 .

Theorem 3.10. *The functor $\mathfrak{A}_{\text{bos}} : \text{GlobHypGreen} \longrightarrow \text{C}^* \text{Alg}$ is a bosonic locally covariant quantum field theory, i.e., the following axioms hold:*

- (i) (**Quantum causality**) *Let (M_j, S_j, P_j) be objects in GlobHypGreen , $j = 1, 2, 3$, and (f_j, F_j) morphisms from (M_j, S_j, P_j) to (M_3, S_3, P_3) , $j = 1, 2$, such that $f_1(M_1)$ and $f_2(M_2)$ are causally disjoint regions in M_3 .*

Then the subalgebras $\mathfrak{A}_{\text{bos}}(f_1, F_1)(\mathfrak{A}_{\text{bos}}(M_1, S_1, P_1))$ and $\mathfrak{A}_{\text{bos}}(f_2, F_2)(\mathfrak{A}_{\text{bos}}(M_2, S_2, P_2))$ of $\mathfrak{A}_{\text{bos}}(M_3, S_3, P_3)$ commute.

- (ii) (**Time slice axiom**) *Let (M_j, S_j, P_j) be objects in GlobHypGreen , $j = 1, 2$, and (f, F) a morphism from (M_1, S_1, P_1) to (M_2, S_2, P_2) such that there is a Cauchy hypersurface $\Sigma \subset M_1$ for which $f(\Sigma)$ is a Cauchy hypersurface of M_2 . Then*

$$\mathfrak{A}_{\text{bos}}(f, F) : \mathfrak{A}_{\text{bos}}(M_1, S_1, P_1) \rightarrow \mathfrak{A}_{\text{bos}}(M_2, S_2, P_2)$$

is an isomorphism.

Proof. We first show (i). For notational simplicity we assume without loss of generality that f_j and F_j are inclusions, $j = 1, 2$. Let $\varphi_j \in C_c^\infty(M_j, S_j)$. Since M_1 and M_2 are causally disjoint, the sections $G\varphi_1$ and φ_2 have disjoint support, thus

$$\omega([\varphi_1], [\varphi_2]) = \int_M \langle G\varphi_1, \varphi_2 \rangle dV = 0.$$

Now relation (iv) in Definition A.11 tells us

$$w([\varphi_1]) \cdot w([\varphi_2]) = w([\varphi_1] + [\varphi_2]) = w([\varphi_2]) \cdot w([\varphi_1]).$$

Since $\mathfrak{A}_{\text{bos}}(f_1, F_1)(\mathfrak{A}_{\text{bos}}(M_1, S_1, P_1))$ is generated by elements of the form $w([\varphi_1])$ and $\mathfrak{A}_{\text{bos}}(f_2, F_2)(\mathfrak{A}_{\text{bos}}(M_2, S_2, P_2))$ by elements of the form $w([\varphi_2])$, the assertion follows.

In order to prove (ii) we show that $\text{SYMPL}(f, F)$ is an isomorphism of symplectic vector spaces provided f maps a Cauchy hypersurface of M_1 onto a Cauchy hypersurface of M_2 . Since symplectic linear maps are always injective, we only need to show surjectivity of $\text{SYMPL}(f, F)$. This is most easily seen by replacing $\text{SYMPL}(M_j, S_j, P_j)$ by $\ker(P_j) \cap C_{\text{sc}}^\infty(M_j, S_j)$ as in Remark 3.9. Again we assume without loss of generality that f and F are inclusions.

Let $\psi \in C_{\text{sc}}^\infty(M_2, S_2)$ be a solution of $P_2\psi = 0$. Let φ be the restriction of ψ to M_1 . Then φ solves $P_1\varphi = 0$ and has spacelike compact support in M_1 by Lemma 3.11 below. We will show that there is only one solution in M_2 with spacelike compact support extending φ . It will then follow that ψ is the image of φ under the extension map corresponding to $\text{SYMPL}(f, F)$ and surjectivity will be shown.

To prove uniqueness of the extension, we may, by linearity, assume that $\varphi = 0$. Then ψ_+ defined by

$$\psi_+(x) := \begin{cases} \psi(x), & \text{if } x \in J_+^{M_2}(\Sigma), \\ 0, & \text{otherwise,} \end{cases}$$

is smooth since ψ vanishes in an open neighborhood of Σ . Now ψ_+ solves $P_2\psi_+ = 0$ and has past-compact support. By Remark 3.6, $\psi_+ \equiv 0$, i.e., ψ vanishes on $J_+^{M_2}(\Sigma)$. One shows similarly that ψ vanishes on $J_-^{M_2}(\Sigma)$, hence $\psi = 0$. \square

Lemma 3.11. *Let M be a globally hyperbolic spacetime and let $M' \subset M$ be a causally compatible open subset which contains a Cauchy hypersurface of M . Let $A \subset M$ be spacelike compact in M .*

Then $A \cap M'$ is spacelike compact in M' .

Proof. Fix a common Cauchy hypersurface Σ of M' and M . By assumption, there exists a compact subset $K \subset M$ with $A \subset J^M(K)$. Then $K' := J^M(K) \cap \Sigma$ is compact [4, Cor. A.5.4] and contained in M' .

Moreover $A \subset J^M(K')$: let $p \in A$ and let γ be a causal curve (in M) from p to some $k \in K$. Then γ can be extended to an inextendible causal curve in M , which hence meets Σ at some point q . Because of $q \in \Sigma \cap J^M(k) \subset K'$ one has $p \in J^M(K')$.

Therefore $A \cap M' \subset J^M(K') \cap M' = J^{M'}(K')$ because of the causal compatibility of M' in M . The lemma is proved. \square

The quantization process described in this subsection applies in particular to formally self-adjoint wave and Dirac-type operators.

3.2. Fermionic quantization. Next we construct a fermionic quantization. For this we need a functorial construction of Hilbert spaces rather than symplectic vector spaces. As we shall see this seems to be possible only under much more restrictive assumptions. The underlying Lorentzian manifold M is assumed to be a globally hyperbolic spacetime as before. The vector bundle S is assumed to be complex with Hermitian inner product $\langle \cdot, \cdot \rangle$ which may be indefinite. The formally self-adjoint Green-hyperbolic operator P is assumed to be of first order.

Definition 3.12. A formally self-adjoint Green-hyperbolic operator P of first order acting on sections of a complex vector bundle S over a spacetime M is of *definite*

type if and only if for any $x \in M$ and any future-directed timelike tangent vector $\mathbf{n} \in T_x M$, the bilinear map

$$S_x \times S_x \rightarrow \mathbb{C}, \quad (\varphi, \psi) \mapsto \langle i\sigma_P(\mathbf{n}^\flat) \cdot \varphi, \psi \rangle,$$

yields a positive definite Hermitian scalar product on S_x .

Example 3.13. The classical Dirac operator P from Example 2.21 is, when defined with the correct sign, of definite type, see e.g. [5, Sec. 1.1.5] or [3, Sec. 2].

Example 3.14. If $E \rightarrow M$ is a semi-Riemannian or -Hermitian vector bundle endowed with a metric connection over a spin spacetime M , then the twisted Dirac operator from Example 2.22 is of definite type if and only if the metric on E is positive definite. This can be seen by evaluating the tensorized inner product on elements of the form $\sigma \otimes v$, where $v \in E_x$ is null.

Example 3.15. The operator $P = i(d - \delta)$ on $S = \Lambda T^*M \otimes \mathbb{C}$ is of Dirac type but not of definite type. This follows from Example 3.14 applied to Example 2.23, since the natural inner product on ΣM is not positive definite. An alternative elementary proof is the following: for any timelike tangent vector \mathbf{n} on M and the corresponding covector \mathbf{n}^\flat , one has

$$\langle i\sigma_P(\mathbf{n}^\flat)\mathbf{n}^\flat, \mathbf{n}^\flat \rangle = -\langle \mathbf{n}^\flat \wedge \mathbf{n}^\flat - \mathbf{n} \lrcorner \mathbf{n}^\flat, \mathbf{n}^\flat \rangle = \langle \mathbf{n}, \mathbf{n} \rangle \langle 1, \mathbf{n}^\flat \rangle = 0.$$

Example 3.16. The Rarita-Schwinger operator defined in Section 2.6 is not of definite type if the dimension of the manifolds is $m \geq 3$. This can be seen as follows. Fix a point $x \in M$ and a pointwise orthonormal basis $(e_j)_{1 \leq j \leq m}$ of $T_x M$ with e_1 timelike. The Lorentzian metric induces inner products on ΣM and on $\Sigma^{3/2}M$ which we denote by $\langle \cdot, \cdot \rangle$. Choose $\xi := e_1^\flat \in T_x^*M$ and $\psi \in \Sigma_x^{3/2}M$. Since $\sigma_{\mathcal{D}}(\xi)$ is pointwise obtained as the orthogonal projection of $\sigma_{\mathcal{D}}(\xi)$ onto $\Sigma_x^{3/2}M$, one has

$$\begin{aligned} \langle -i\sigma_{\mathcal{D}}(\xi)\psi, \psi \rangle &= \langle (\text{id} \otimes \xi^\sharp \cdot)\psi, \psi \rangle - \underbrace{\frac{2}{m} \sum_{\beta=1}^m \langle e_\beta^* \otimes e_\beta \cdot \psi_1, \psi \rangle}_{=0} \\ &= \sum_{\beta=1}^m \varepsilon_\beta \langle e_1 \cdot \psi_\beta, \psi_\beta \rangle. \end{aligned}$$

Choose, as in the proof of Lemma 2.26, a $\psi \in \Sigma_x^{3/2}M$ with $\psi_k = 0$ for all $3 \leq k \leq m$. For such a ψ the condition $\psi \in \Sigma_x^{3/2}M$ becomes $e_1 \cdot \psi_1 = e_2 \cdot \psi_2$. As in the proof of Lemma 2.26 we obtain

$$\langle -i\sigma_{\mathcal{D}}(\xi)\psi, \psi \rangle = -\langle e_1 \cdot \psi_2, \psi_2 \rangle + \langle e_1 \cdot \psi_2, \psi_2 \rangle = 0,$$

which shows that the Rarita-Schwinger operator cannot be of definite type.

We define the category GlobHypDef , whose objects are the triples (M, S, P) , where M is a globally hyperbolic spacetime, S is a complex vector bundle equipped with a complex inner product $\langle \cdot, \cdot \rangle$, and P is a formally self-adjoint Green-hyperbolic operator of definite type acting on sections of S . The morphisms are the same as in the category GlobHypGreen .

We construct a covariant functor from GlobHypDef to HILB , where HILB denotes the category whose objects are complex pre-Hilbert spaces and whose morphisms are isometric linear embeddings. As in Section 3.1, the underlying vector space

is the space of classical solutions to the equation $P\varphi = 0$ with spacelike compact support. We put

$$\text{SOL}(M, S, P) := \ker(P) \cap C_{\text{sc}}^\infty(M, S).$$

Here ‘‘SOL’’ stands for classical solutions of the equation $P\varphi = 0$ with spacelike compact support.

Lemma 3.17. *Let (M, S, P) be an object in GlobHypDef. Let $\Sigma \subset M$ be a smooth spacelike Cauchy hypersurface with its future-oriented unit normal vector field \mathbf{n} and its induced volume element dA . Then*

$$(8) \quad (\varphi, \psi) := \int_{\Sigma} \langle i\sigma_P(\mathbf{n}^\flat) \cdot \varphi|_{\Sigma}, \psi|_{\Sigma} \rangle dA,$$

yields a positive definite Hermitian scalar product on $\text{SOL}(M, S, P)$ which does not depend on the choice of Σ .

Proof. First note that $\text{supp}(\varphi) \cap \Sigma$ is compact since $\text{supp}(\varphi)$ is spacelike compact, so that the integral is well-defined. We have to show that it does not depend on the choice of Cauchy hypersurface. Let Σ' be any other smooth spacelike Cauchy hypersurface. Assume first that Σ and Σ' are disjoint and let Ω be the domain enclosed by Σ and Σ' in M . Its boundary is $\partial\Omega = \Sigma \cup \Sigma'$. Without loss of generality, one may assume that $\Sigma' \subset J_+^M(\Sigma)$. By the Green’s formula [40, p. 160, Prop. 9.1] we have for all $\varphi, \psi \in C_{\text{sc}}^\infty(M, S)$,

$$(9) \quad \int_{\Omega} (\langle P\varphi, \psi \rangle - \langle \varphi, P\psi \rangle) dV = \int_{\Sigma'} \langle \sigma_P(\mathbf{n}^\flat)\varphi, \psi \rangle dA - \int_{\Sigma} \langle \sigma_P(\mathbf{n}^\flat)\varphi, \psi \rangle dA.$$

For $\varphi, \psi \in \text{SOL}(M, S, P)$ we have $P\varphi = P\psi = 0$ and thus

$$0 = \int_{\Sigma} \langle \sigma_P(\mathbf{n}^\flat)\varphi, \psi \rangle dA - \int_{\Sigma'} \langle \sigma_P(\mathbf{n}^\flat)\varphi, \psi \rangle dA.$$

This shows the result in the case $\Sigma \cap \Sigma' = \emptyset$.

If $\Sigma \cap \Sigma' \neq \emptyset$ consider the subset $I_+^M(\Sigma) \cap I_-^M(\Sigma')$ of M where, as usual, $I_+^M(\Sigma)$ and $I_-^M(\Sigma)$ denote the chronological future and past of the subset Σ in M , respectively. This subset is nonempty, open, and globally hyperbolic. This follows e.g. from [4, Lemma A.5.8]. Hence it admits a smooth spacelike Cauchy hypersurface Σ'' by Theorem 2.3. By construction, Σ'' meets neither Σ nor Σ' and it can be easily checked that Σ'' is also a Cauchy hypersurface of M . The result follows from the argument above being applied first to the pair (Σ, Σ'') and then to the pair (Σ'', Σ') . \square

Remark 3.18. If one drops the assumption that P be of definite type, then the above sesquilinear form (\cdot, \cdot) on $\ker(P) \cap C_{\text{sc}}^\infty(M, S)$ still does not depend on the choice of Σ , however it need no longer be positive definite and can even be degenerate. Pick for instance the spin Dirac operator D_g associated to the underlying Lorentzian metric g on a spin spacetime M (see Example 2.21) and, keeping the spinor bundle $\Sigma_g M$ associated to g , change the metric on M so that the new metric g' has larger future and past cones at each point. Note that this implies that any globally hyperbolic subregion of (M, g') is also globally hyperbolic in (M, g) . Then, denoting by D_g^* the formal adjoint of D_g with respect to the metric g' , the operator $\begin{pmatrix} 0 & D_g \\ D_g^* & 0 \end{pmatrix}$ on $\Sigma_g M \oplus \Sigma_g M$ remains Green-hyperbolic but it fails to be of definite type, since there exist timelike vectors for g' which are lightlike for g .

Hence the principal symbol of the operator becomes non-invertible and the bilinear form in (8) becomes degenerate for these g' -timelike covectors.

For any object (M, S, P) in GlobHypDef we will from now on equip $\text{SOL}(M, S, P)$ with the Hermitian scalar product in (8) and thus turn $\text{SOL}(M, S, P)$ into a pre-Hilbert space.

Given a morphism (f, F) from (M_1, S_1, P_1) to (M_2, S_2, P_2) in GlobHypDef , then this is also a morphism in GlobHypGreen and hence induces a homomorphism $\text{SYMPL}(f, F) : \text{SYMPL}(M_1, S_1, P_1) \rightarrow \text{SYMPL}(M_2, S_2, P_2)$. As explained in Remark 3.9, there is a corresponding extension homomorphism $\text{SOL}(f, F) : \text{SOL}(M_1, S_1, P_1) \rightarrow \text{SOL}(M_2, S_2, P_2)$. In other words, $\text{SOL}(f, F)$ is defined such that the diagram

$$(10) \quad \begin{array}{ccc} \text{SYMPL}(M_1, S_1, P_1) & \xrightarrow{\text{SYMPL}(f, F)} & \text{SYMPL}(M_2, S_2, P_2) \\ \cong \downarrow & & \downarrow \cong \\ \text{SOL}(M_1, S_1, P_1) & \xrightarrow{\text{SOL}(f, F)} & \text{SOL}(M_2, S_2, P_2) \end{array}$$

commutes. The vertical arrows are the vector space isomorphisms induced by the Green's propagators G_1 and G_2 , respectively.

Lemma 3.19. *The vector space homomorphism $\text{SOL}(f, F) : \text{SOL}(M_1, S_1, P_1) \rightarrow \text{SOL}(M_2, S_2, P_2)$ preserves the scalar products, i.e., it is an isometric linear embedding of pre-Hilbert spaces.*

Proof. Without loss of generality we assume that f and F are inclusions. Let Σ_1 be a spacelike Cauchy hypersurface of M_1 . Let $\varphi_1, \psi_1 \in C_{\text{sc}}^\infty(M_1, S_1)$. Denote the extension of φ_1 by $\varphi_2 := \text{SOL}(f, F)(\varphi_1)$ and similarly for ψ_1 .

Let $K_1 \subset M_1$ be a compact subset such that $\text{supp}(\varphi_2) \subset J^{M_2}(K_1)$ and $\text{supp}(\psi_2) \subset J^{M_2}(K_1)$. We choose a compact submanifold $K \subset \Sigma_1$ with boundary such that $J^{M_1}(K_1) \cap \Sigma_1 \subset K$. Since Σ_1 is a Cauchy hypersurface in M_1 , $J^{M_1}(K_1) \subset J^{M_1}(J^{M_1}(K_1) \cap \Sigma_1) \subset J^{M_1}(K)$.

By Theorem 2.5 there is a spacelike Cauchy hypersurface $\Sigma_2 \subset M_2$ containing K . Since Σ_i is a Cauchy hypersurface of M_i (where $i = 1, 2$), it is met by every inextendible causal curve [30, Lemma 14.29]. Moreover, by definition of a Cauchy hypersurface, Σ_i is achronal in M_i . Since it is also spacelike, Σ_i is even acausal [30, Lemma 14.42]. In particular, it is met *exactly once* by every inextendible causal curve in M_i .

This implies $J^{M_2}(K_1) \subset J^{M_2}(K)$: namely, pick $p \in J^{M_2}(K_1)$ and a causal curve γ in M_2 from p to some $k_1 \in K_1$. Extend γ to an inextendible causal curve $\bar{\gamma}$ in M_2 . Then $\bar{\gamma}$ meets Σ_2 at some point q_2 , because Σ_2 is a Cauchy hypersurface in M_2 . But $\bar{\gamma} \cap M_1$ is also an inextendible causal curve in M_1 , hence it intersects Σ_1 at a point q_1 , which must lie in K by definition of K . Because of $K \subset \Sigma_2$ and the uniqueness of the intersection point, one has $q_1 = q_2$. In particular, $p \in J^{M_2}(K)$.




Fig. 1

We conclude $\text{supp}(\varphi_2) \subset J^{M_2}(K)$. Since $K \subset \Sigma_2$, we have $\text{supp}(\varphi_2) \cap \Sigma_2 \subset J^{M_2}(K) \cap \Sigma_2$ and $J^{M_2}(K) \cap \Sigma_2 = K$ using the acausality of Σ_2 . This shows $\text{supp}(\varphi_2) \cap \Sigma_2 = \text{supp}(\varphi_1) \cap \Sigma_1$ and similarly for ψ_2 . Now we get

$$(\varphi_2, \psi_2) = \int_{\Sigma_2} \langle i\sigma_{P_2}(\mathbf{n}^\flat) \cdot \varphi_2, \psi_2 \rangle dA = \int_{\Sigma_1} \langle i\sigma_{P_1}(\mathbf{n}^\flat) \cdot \varphi_1, \psi_1 \rangle dA = (\varphi_1, \psi_1)$$

and the lemma is proved. \square

The functoriality of SYMPL and diagram (10) show that SOL is a functor from GlobHypDef to HILB, the category of complex pre-Hilbert spaces with isometric linear embeddings. Composing with the functor CAR (see Section A.1), we obtain the covariant functor

$$\mathfrak{A}_{\text{ferm}} := \text{CAR} \circ \text{SOL} : \text{GlobHypDef} \longrightarrow \text{C}^* \text{Alg}.$$

The fermionic algebras $\mathfrak{A}_{\text{ferm}}(M, S, P)$ are actually \mathbb{Z}_2 -graded algebras, see Proposition A.5 (iii).

Theorem 3.20. *The functor $\mathfrak{A}_{\text{ferm}} : \text{GlobHypDef} \longrightarrow \text{C}^* \text{Alg}$ is a fermionic locally covariant quantum field theory, i.e., the following axioms hold:*

- (i) **(Quantum causality)** Let (M_j, S_j, P_j) be objects in GlobHypDef, $j = 1, 2, 3$, and (f_j, F_j) morphisms from (M_j, S_j, P_j) to (M_3, S_3, P_3) , $j = 1, 2$, such that $f_1(M_1)$ and $f_2(M_2)$ are causally disjoint regions in M_3 . Then the subalgebras $\mathfrak{A}_{\text{ferm}}(f_1, F_1)(\mathfrak{A}_{\text{ferm}}(M_1, S_1, P_1))$ and $\mathfrak{A}_{\text{ferm}}(f_2, F_2)(\mathfrak{A}_{\text{ferm}}(M_2, S_2, P_2))$ of $\mathfrak{A}_{\text{ferm}}(M_3, S_3, P_3)$ super-commute¹.
- (ii) **(Time slice axiom)** Let (M_j, S_j, P_j) be objects in GlobHypDef, $j = 1, 2$, and (f, F) a morphism from (M_1, S_1, P_1) to (M_2, S_2, P_2) such that there is a Cauchy hypersurface $\Sigma \subset M_1$ for which $f(\Sigma)$ is a Cauchy hypersurface of M_2 . Then

$$\mathfrak{A}_{\text{ferm}}(f, F) : \mathfrak{A}_{\text{ferm}}(M_1, S_1, P_1) \rightarrow \mathfrak{A}_{\text{ferm}}(M_2, S_2, P_2)$$

is an isomorphism.

¹This means that the odd parts of the algebras anti-commute while the even parts commute with everything.

Proof. To show (i), we assume without loss of generality that f_j and F_j are inclusions. Let $\varphi_1 \in \text{SOL}(M_1, S_1, P_1)$ and $\psi_1 \in \text{SOL}(M_2, S_2, P_2)$. Denote the extensions to M_3 by $\varphi_2 := \text{SOL}(f_1, F_1)(\varphi_1)$ and $\psi_2 := \text{SOL}(f_2, F_2)(\psi_1)$. Choose a compact submanifold K_1 (with boundary) in a spacelike Cauchy hypersurface Σ_1 of M_1 such that $\text{supp}(\varphi_1) \cap \Sigma_1 \subset K_1$ and similarly K_2 for ψ_1 . Since M_1 and M_2 are causally disjoint, $K_1 \cup K_2$ is acausal. Hence, by Theorem 2.5, there exists a Cauchy hypersurface Σ_3 of M_3 containing K_1 and K_2 . As in the proof of Lemma 3.19 one sees that $\text{supp}(\varphi_2) \cap \Sigma_3 = \text{supp}(\varphi_1) \cap \Sigma_1$ and similarly for ψ_2 . Thus, when restricted to Σ_3 , φ_2 and ψ_2 have disjoint support. Hence $(\varphi_2, \psi_2) = 0$. This shows that the subspaces $\text{SOL}(f_1, F_1)(\text{SOL}(M_1, S_1, P_1))$ and $\text{SOL}(f_2, F_2)(\text{SOL}(M_2, S_2, P_2))$ of $\text{SOL}(M_3, S_3, P_3)$ are perpendicular. Definition A.1 shows that the corresponding CAR-algebras must super-commute.

To see (ii) we recall that (f, F) is also a morphism in GlobHypGreen and that we know from Theorem 3.10 that $\text{SYMPL}(f, F)$ is an isomorphism. From diagram (10) we see that $\text{SOL}(f, F)$ is an isomorphism. Hence $\mathfrak{A}_{\text{ferm}}(f, F)$ is also an isomorphism. \square

Remark 3.21. Since causally disjoint regions should lead to commuting observables also in the fermionic case, one usually considers only the even part $\mathfrak{A}_{\text{ferm}}^{\text{even}}(M, S, P)$ (or a subalgebra thereof) as the observable algebra while the full algebra $\mathfrak{A}_{\text{ferm}}(M, S, P)$ is called the *field algebra*.

There is a slightly different description of the functor $\mathfrak{A}_{\text{ferm}}$. Let $\text{HILB}_{\mathbb{R}}$ denote the category whose objects are the real pre-Hilbert spaces and whose morphisms are the isometric linear embeddings. We have the functor $\text{REAL} : \text{HILB} \rightarrow \text{HILB}_{\mathbb{R}}$ which associates to each complex pre-Hilbert space $(V, (\cdot, \cdot))$ its underlying real pre-Hilbert space $(V, \Re(\cdot, \cdot))$. By Remark A.10,

$$\mathfrak{A}_{\text{ferm}} = \text{CAR}_{\text{sd}} \circ \text{REAL} \circ \text{SOL}.$$

Since the self-dual CAR-algebra of a real pre-Hilbert space is the Clifford algebra of its complexification and since for any complex pre-Hilbert space V we have

$$\text{REAL}(V) \otimes_{\mathbb{R}} \mathbb{C} = V \oplus V^*,$$

$\mathfrak{A}_{\text{ferm}}(M, S, P)$ is also the Clifford algebra of $\text{SOL}(M, S, P) \oplus \text{SOL}(M, S, P)^* = \text{SOL}(M, S \oplus S^*, P \oplus P^*)$. This is the way this functor is often described in the physics literature, see e.g. [39, p. 115f].

Self-dual CAR-representations are more natural for real fields. Let M be globally hyperbolic and let $S \rightarrow M$ be a *real* vector bundle equipped with a real inner product $\langle \cdot, \cdot \rangle$. A formally skew-adjoint² differential operator P acting on sections of S is called of *definite type* if and only if for any $x \in M$ and any future-directed timelike tangent vector $\mathfrak{n} \in T_x M$, the bilinear map

$$S_x \times S_x \rightarrow \mathbb{R}, \quad (\varphi, \psi) \mapsto \langle \sigma_P(\mathfrak{n}^b) \cdot \varphi, \psi \rangle,$$

yields a positive definite Euclidean scalar product on S_x . An example is given by the real Dirac operator

$$D := \sum_{j=1}^m \varepsilon_j e_j \cdot \nabla_{e_j}$$

acting on sections of the real spinor bundle $\Sigma^{\mathbb{R}} M$.

²instead of self-adjoint!

Given a smooth spacelike Cauchy hypersurface $\Sigma \subset M$ with future-directed time-like unit normal field \mathbf{n} , we define a scalar product on $\text{SOL}(M, S, P) = \ker(P) \cap C_{\text{sc}}^\infty(M, S, P)$ by

$$(\varphi, \psi) := \int_{\Sigma} \langle \sigma_P(\mathbf{n}^b) \cdot \varphi|_{\Sigma}, \psi|_{\Sigma} \rangle dA.$$

With essentially the same proofs as before, one sees that this scalar product does not depend on the choice of Cauchy hypersurface Σ and that a morphism $(f, F) : (M_1, S_1, P_1) \rightarrow (M_2, S_2, P_2)$ gives rise to an extension operator $\text{SOL}(f, F) : \text{SOL}(M_1, S_1, P_1) \rightarrow \text{SOL}(M_2, S_2, P_2)$ preserving the scalar product. We have constructed a functor

$$\text{SOL} : \text{GlobHypSkewDef} \longrightarrow \text{HILB}_{\mathbb{R}}$$

where GlobHypSkewDef denotes the category whose objects are triples (M, S, P) with M globally hyperbolic, $S \rightarrow M$ a real vector bundle with real inner product and P a formally skew-adjoint, Green-hyperbolic differential operator of definite type acting on sections of S . The morphisms are the same as before.

Now the functor

$$\mathfrak{A}_{\text{ferm}}^{\text{sd}} := \text{CAR}_{\text{sd}} \circ \text{SOL} : \text{GlobHypSkewDef} \longrightarrow \text{C}^* \text{Alg}$$

is a locally covariant quantum field theory in the sense that Theorem 3.20 holds with $\mathfrak{A}_{\text{ferm}}$ replaced by $\mathfrak{A}_{\text{ferm}}^{\text{sd}}$.

4. STATES AND QUANTUM FIELDS

In order to produce numbers out of our quantum field theory that can be compared to experiments, we need states, in addition to observables. We briefly recall the relation between states and representations via the GNS-construction. Then we show how the choice of a state gives rise to quantum fields and n -point functions.

4.1. States and representations. Recall that a *state* on a unital C^* -algebra A is a linear functional $\tau : A \rightarrow \mathbb{C}$ such that

- (i) τ is positive, i.e., $\tau(a^*a) \geq 0$ for all $a \in A$;
- (ii) τ is normed, i.e., $\tau(1) = 1$.

One checks that for any state the sesquilinear form $A \times A \rightarrow \mathbb{C}$, $(a, b) \mapsto \tau(b^*a)$, is a positive semi-definite Hermitian product and $|\tau(a)| \leq \|a\|$ for all $a \in A$. In particular, τ is continuous.

Any state induces a representation of A . Namely, the sesquilinear form $\tau(b^*a)$ induces a scalar product $\langle \cdot, \cdot \rangle_{\tau}$ on $A/\{a \in A \mid \tau(a^*a) = 0\}$. The Hilbert space completion of $A/\{a \in A \mid \tau(a^*a) = 0\}$ is denoted by \mathcal{H}_{τ} . The action of A on \mathcal{H}_{τ} is induced by the multiplication in A ,

$$\pi_{\tau}(a)[b]_{\tau} := [ab]_{\tau},$$

where $[a]_{\tau}$ denotes the residue class of $a \in A$ in $A/\{a \in A \mid \tau(a^*a) = 0\}$. This representation is known as the *GNS-representation* induced by τ . The residue class $\Omega_{\tau} := [1]_{\tau} \in \mathcal{H}_{\tau}$ is called the *vacuum vector*. By construction, it is a cyclic vector, i.e., the orbit $\pi_{\tau}(A) \cdot \Omega_{\tau} = A/\{a \in A \mid \tau(a^*a) = 0\}$ is dense in \mathcal{H}_{τ} .

The GNS-representation together with the vacuum vector allows to reconstruct the state since

$$(11) \quad \tau(a) = \tau(1^*a1) = \langle \pi_{\tau}(a)\Omega_{\tau}, \Omega_{\tau} \rangle_{\tau}.$$

If we look at the vector state $\tilde{\tau} : \mathcal{L}(\mathcal{H}_\tau) \rightarrow \mathbb{C}$, $\tilde{\tau}(\tilde{a}) = \langle \tilde{a}\Omega_\tau, \Omega_\tau \rangle_\tau$, on the C^* -algebra $\mathcal{L}(\mathcal{H}_\tau)$ of bounded linear operators on \mathcal{H}_τ , then (11) says that the diagram

$$\begin{array}{ccc} A & \xrightarrow{\pi_\tau} & \mathcal{L}(\mathcal{H}_\tau) \\ & \searrow \tau & \swarrow \tilde{\tau} \\ & & \mathbb{C} \end{array}$$

commutes. One checks that $\|\pi_\tau\| \leq 1$, see [2, p. 20]. In particular, $\pi_\tau : A \rightarrow \mathcal{L}(\mathcal{H}_\tau)$ is continuous.

See e.g. [2, Sec. 1.4] or [9, Sec. 2.3] for details on states and representations of C^* -algebras.

4.2. Bosonic quantum field. Now let (M, S, P) be an object in GlobHypGreen and τ a state on the corresponding bosonic algebra $\mathfrak{A}_{\text{bos}}(M, S, P)$. Intuitively, the quantum field should be an operator-valued distribution Φ on M such that

$$e^{i\Phi(f)} = w([f])$$

for all test sections $f \in C_c^\infty(M, S)$. Here $[f]$ denotes the residue class in $\text{SYMPL}(M, S, P) = C_c^\infty(M, S) / \ker G$ and $w : \text{SYMPL}(M, S, P) \rightarrow \mathfrak{A}_{\text{bos}}(M, S, P)$ is as in Definition A.11. This suggests the definition

$$\Phi(f) := -i \left. \frac{d}{dt} \right|_{t=0} w(t[f]).$$

The problem is that w is highly discontinuous so that this derivative does not make sense. This is where states and representations come into the play. We call a state τ on $\mathfrak{A}_{\text{bos}}(M, S, P)$ *regular* if for each $f \in C_c^\infty(M, S)$ and each $h \in \mathcal{H}_\tau$ the map $t \mapsto \pi_\tau(w(t[f]))h$ is continuous. Then $t \mapsto \pi_\tau(w(t[f]))$ is a strongly continuous one-parameter unitary group for any $f \in C_c^\infty(M, S)$ because

$$\pi_\tau(w((t+s)[f])) = \pi_\tau(e^{i\omega(t[f], s[f])/2} w(t[f])w(s[f])) = \pi_\tau(w(t[f]))\pi_\tau(w(s[f])).$$

Here we used Definition A.11 (iv) and the fact that ω is skew-symmetric so that $\omega(t[f], s[f]) = 0$. By Stone's theorem [34, Thm. VIII.8] this one-parameter group has a unique infinitesimal generator, i.e., a self-adjoint, generally unbounded operator $\Phi_\tau(f)$ on \mathcal{H}_τ such that

$$e^{it\Phi_\tau(f)} = \pi_\tau(w(t[f])).$$

For all h in the domain of $\Phi_\tau(f)$ we have

$$\Phi_\tau(f)h = -i \left. \frac{d}{dt} \right|_{t=0} \pi_\tau(w(t[f]))h.$$

We call the operator-valued map $f \mapsto \Phi_\tau(f)$ the *quantum field* corresponding to τ .

Definition 4.1. A regular state τ on $\mathfrak{A}_{\text{bos}}(M, S, P)$ is called *strongly regular* if

- (i) there is a dense subspace $\mathcal{D}_\tau \subset \mathcal{H}_\tau$ contained in the domain of $\Phi_\tau(f)$ for any $f \in C_c^\infty(M, S)$;
- (ii) $\Phi_\tau(f)(\mathcal{D}_\tau) \subset \mathcal{D}_\tau$ for any $f \in C_c^\infty(M, S)$;
- (iii) the map $C_c^\infty(M, S) \rightarrow \mathcal{H}_\tau$, $f \mapsto \Phi_\tau(f)h$, is continuous for every fixed $h \in \mathcal{D}_\tau$.

For a strongly regular state τ we have for all $f, g \in C_c^\infty(M, S)$, $\alpha, \beta \in \mathbb{R}$ and $h \in \mathcal{D}_\tau$:

$$\begin{aligned} \Phi_\tau(\alpha f + \beta g)h &= -i \frac{d}{dt} \Big|_{t=0} \pi_\tau(w(t[\alpha f + \beta g]))h \\ &= -i \frac{d}{dt} \Big|_{t=0} \left\{ e^{i\alpha\beta t^2 \omega([f],[g])/2} \pi_\tau(w(\alpha t[f])) \pi_\tau(w(\beta t[g]))h \right\} \\ &= -i \frac{d}{dt} \Big|_{t=0} \pi_\tau(w(\alpha t[f]))h - i \frac{d}{dt} \Big|_{t=0} \pi_\tau(w(\beta t[g]))h \\ &= \alpha \Phi_\tau(f)h + \beta \Phi_\tau(g)h. \end{aligned}$$

Hence $\Phi_\tau(f)$ depends linearly on f . The quantum field Φ_τ is therefore a distribution on M with values in self-adjoint operators on \mathcal{H}_τ .

The n -point functions are defined by

$$\begin{aligned} \tau_n(f_1, \dots, f_n) &:= \langle \Phi_\tau(f_1) \cdots \Phi_\tau(f_n) \Omega_\tau, \Omega_\tau \rangle_\tau \\ &= \tilde{\tau}(\Phi_\tau(f_1) \cdots \Phi_\tau(f_n)) \\ &= \tilde{\tau} \left(\left(-i \frac{d}{dt_1} \Big|_{t_1=0} \pi_\tau(w(t_1[f_1])) \right) \cdots \left(-i \frac{d}{dt_n} \Big|_{t_n=0} \pi_\tau(w(t_n[f_n])) \right) \right) \\ &= (-i)^n \frac{\partial^n}{\partial t_1 \cdots \partial t_n} \Big|_{t_1=\dots=t_n=0} \tilde{\tau}(\pi_\tau(w(t_1[f_1])) \cdots \pi_\tau(w(t_n[f_n]))) \\ &= (-i)^n \frac{\partial^n}{\partial t_1 \cdots \partial t_n} \Big|_{t_1=\dots=t_n=0} \tilde{\tau}(\pi_\tau(w(t_1[f_1]) \cdots w(t_n[f_n]))) \\ &= (-i)^n \frac{\partial^n}{\partial t_1 \cdots \partial t_n} \Big|_{t_1=\dots=t_n=0} \tau(w(t_1[f_1]) \cdots w(t_n[f_n])). \end{aligned}$$

For a strongly regular state τ the n -point functions are continuous separately in each factor. By the Schwartz kernel theorem [23, Thm. 5.2.1] the n -point function τ_n extends uniquely to a distribution on $M \times \cdots \times M$ (n times) in the following sense: Let $S^* \boxtimes \cdots \boxtimes S^*$ be the bundle over $M \times \cdots \times M$ whose fiber over (x_1, \dots, x_n) is given by $S_{x_1}^* \otimes \cdots \otimes S_{x_n}^*$. Then there is a unique distribution on $M \times \cdots \times M$ in the bundle $S^* \boxtimes \cdots \boxtimes S^*$, again denoted τ_n , such that for all $f_j \in C_c^\infty(M, S)$,

$$\tau_n(f_1, \dots, f_n) = \tau_n(f_1 \otimes \cdots \otimes f_n)$$

where $(f_1 \otimes \cdots \otimes f_n)(x_1, \dots, x_n) := f_1(x_1) \otimes \cdots \otimes f_n(x_n)$.

Theorem 4.2. *Let (M, S, P) be an object in GlobHypGreen and τ a strongly regular state on the corresponding bosonic algebra $\mathfrak{A}_{\text{bos}}(M, S, P)$. Then*

- (i) $P\Phi_\tau = 0$ and $P\tau_n(f_1, \dots, f_{j-1}, \cdot, f_{j+1}, \dots, f_n) = 0$ hold in the distributional sense where $f_k \in C_c^\infty(M, S)$, $k \neq j$, are fixed;
- (ii) the quantum field satisfies the canonical commutation relations, i.e.,

$$[\Phi_\tau(f), \Phi_\tau(g)]h = i \int_M \langle Gf, g \rangle dV \cdot h$$

for all $f, g \in C_c^\infty(M, S)$ and $h \in \mathcal{D}_\tau$;

(iii) the n -point functions satisfy the canonical commutation relations, i.e.,

$$\begin{aligned} & \tau_{n+2}(f_1, \dots, f_{j-1}, f_j, f_{j+1}, \dots, f_{n+2}) \\ & - \tau_{n+2}(f_1, \dots, f_{j-1}, f_{j+1}, f_j, f_{j+2}, \dots, f_{n+2}) \\ & = i \int_M \langle Gf_j, f_{j+1} \rangle dV \cdot \tau_n(f_1, \dots, f_{j-1}, f_{j+2}, \dots, f_{n+2}) \end{aligned}$$

for all $f_1, \dots, f_{n+2} \in C_c^\infty(M, S)$.

Proof. Since P is formally self-adjoint and $GPf = 0$ for any $f \in C_c^\infty(M, S)$, we have for any $h \in \mathcal{D}_\tau$:

$$(P\Phi_\tau)(f)h = \Phi_\tau(Pf)h = -i \left. \frac{d}{dt} \right|_{t=0} \pi_\tau(\underbrace{w(t[Pf])}_{=0})h = -i \left. \frac{d}{dt} \right|_{t=0} h = 0.$$

This shows $P\Phi_\tau = 0$. The result for the n -point functions follows and (i) is proved. To show (ii) we observe that by Definition A.11 (iv) we have on the one hand

$$w([f + g]) = e^{i\omega([f],[g])/2} w([f])w([g])$$

and on the other hand

$$w([f + g]) = e^{i\omega([g],[f])/2} w([g])w([f]),$$

hence

$$w([f])w([g]) = e^{-i\omega([f],[g])} w([g])w([f]).$$

Thus

$$\begin{aligned} \Phi_\tau(f)\Phi_\tau(g)h &= - \left. \frac{\partial^2}{\partial t \partial s} \right|_{t=s=0} \pi_\tau(w(t[f])w(s[g]))h \\ &= - \left. \frac{\partial^2}{\partial t \partial s} \right|_{t=s=0} \pi_\tau(e^{-i\omega(t[f],s[g])} w(s[g])w(t[f]))h \\ &= - \left. \frac{\partial^2}{\partial t \partial s} \right|_{t=s=0} \left\{ e^{-i\omega(t[f],s[g])} \cdot \pi_\tau(w(s[g])w(t[f]))h \right\} \\ &= i\omega([f],[g])h + \Phi_\tau(g)\Phi_\tau(f)h \\ &= i \int_M \langle Gf, g \rangle dV \cdot h + \Phi_\tau(g)\Phi_\tau(f)h. \end{aligned}$$

This shows (ii). Assertion (iii) follows from (ii). \square

Remark 4.3. As a consequence of the canonical commutation relations we get

$$[\Phi_\tau(f), \Phi_\tau(g)] = 0$$

if the supports of f and g are causally disjoint, i.e., if there is no causal curve from $\text{supp}(f)$ to $\text{supp}(g)$. The reason is that in this case the supports of Gf and g are disjoint. A similar remark holds for the n -point functions.

Remark 4.4. In the physics literature one also finds the statement $\Phi(\bar{f}) = \Phi(f)^*$. This simply expresses the fact that we are dealing with a theory over the reals. We have encoded this by considering real vector bundles S , see Definition 3.1, and the fact that $\Phi_\tau(f)$ is always self-adjoint.

4.3. Fermionic quantum fields. Let (M, S, P) be an object in GlobHypDef and let τ be a state on the fermionic algebra $\mathfrak{A}_{\text{ferm}}(M, S, P)$. For $f \in C_c^\infty(M, S)$ we put

$$\begin{aligned}\Phi_\tau(f) &:= -\pi_\tau(\mathbf{a}(Gf)^*), \\ \Phi_\tau^+(f) &:= \pi_\tau(\mathbf{a}(Gf)),\end{aligned}$$

where \mathbf{a} is as in Definition A.1 (compare [18, Sec. III.B, p. 141]). Since π_τ , \mathbf{a} , and G are sequentially continuous (for G see [4, Prop. 3.4.8]), so are Φ_τ and Φ_τ^+ . In contrast to the bosonic case, no regularity assumption on τ is needed. Hence Φ_τ and Φ_τ^+ are distributions on M with values in the space of bounded operators on \mathcal{H}_τ . Note that Φ_τ is linear while Φ_τ^+ is anti-linear.

Theorem 4.5. *Let (M, S, P) be an object in GlobHypDef and τ a state on the corresponding fermionic algebra $\mathfrak{A}_{\text{ferm}}(M, S, P)$. Then*

- (i) $P\Phi_\tau = P\Phi_\tau^+ = 0$ holds in the distributional sense;
- (ii) the quantum fields satisfy the canonical anti-commutation relations, i.e.,

$$\begin{aligned}\{\Phi_\tau(f), \Phi_\tau(g)\} &= \{\Phi_\tau^+(f), \Phi_\tau^+(g)\} = 0, \\ \{\Phi_\tau(f), \Phi_\tau^+(g)\} &= i\left(\int_M \langle Gf, g \rangle dV\right) \cdot \text{id}_{\mathcal{H}_\tau}\end{aligned}$$

for all $f, g \in C_c^\infty(M, S)$.

Proof. Since $GP = 0$ on $C_c^\infty(M, S)$, we have $P\Phi_\tau(f) = \Phi_\tau(Pf) = -\pi_\tau(\mathbf{a}(GPf)^*) = 0$ and similarly for Φ_τ^+ . This proves assertion (i).

Using Definition A.1 (ii) we compute

$$\begin{aligned}\{\Phi_\tau(f), \Phi_\tau(g)\} &= \{\pi_\tau(\mathbf{a}(Gf)^*), \pi_\tau(\mathbf{a}(Gg)^*)\} \\ &= \pi_\tau(\{\mathbf{a}(Gf)^*, \mathbf{a}(Gg)^*\}) \\ &= \pi_\tau(\{\mathbf{a}(Gg), \mathbf{a}(Gf)\}^*) \\ &= 0.\end{aligned}$$

Similarly one sees $\{\Phi_\tau^+(f), \Phi_\tau^+(g)\} = 0$. Definition A.1 (iii) also yields

$$\{\Phi_\tau(f), \Phi_\tau^+(g)\} = -\pi_\tau(\{\mathbf{a}(Gf)^*, \mathbf{a}(Gg)\}) = -(Gf, Gg) \cdot \text{id}_{\mathcal{H}_\tau}.$$

To prove assertion (ii) we have to verify

$$(12) \quad (Gf, Gg) = -i \int_M \langle Gf, g \rangle dV$$

Let $\Sigma \subset M$ be a smooth spacelike Cauchy hypersurface. Since $\text{supp}(G_+g)$ is past-compact, we can find a Cauchy hypersurface $\Sigma' \subset M$ in the past of Σ which does not intersect $\text{supp}(G_+g) \subset J_+^M(\text{supp}(g))$. Denote the region between Σ and Σ' by Ω' . The Green's formula (9) yields

$$\begin{aligned}(Gf, G_+g) &= \int_\Sigma \langle i\sigma_P(\mathbf{n}^\flat) \cdot Gf, G_+g \rangle dA \\ &= \int_{\Sigma'} \langle i\sigma_P(\mathbf{n}^\flat) \cdot Gf, G_+g \rangle dA + i \int_{\Omega'} (\langle PGf, G_+g \rangle - \langle Gf, PG_+g \rangle) dV \\ &= -i \int_{\Omega'} \langle Gf, g \rangle dV\end{aligned}$$

because $PG_+g = g$ and $PGf = 0$. Since Σ' can be chosen arbitrarily to the past, this shows

$$(13) \quad (Gf, G_+g) = -i \int_{J_-(\Sigma)} \langle Gf, g \rangle dV.$$

A similar computation yields

$$(14) \quad (Gf, G_-g) = i \int_{J_+(\Sigma)} \langle Gf, g \rangle dV.$$

Subtracting (14) from (13) yields (12) and concludes the proof of assertion (ii). \square

Remark 4.6. Similarly to the bosonic case, we find

$$\{\Phi_\tau(f), \Phi_\tau^+(g)\} = 0$$

if the supports of f and g are causally disjoint.

Remark 4.7. Using the anti-commutation relations in Theorem 4.5 (ii), the computation of n -point functions can be reduced to those of the form

$$\tau_{n,n'}(f_1, \dots, f_n, g_1, \dots, g_{n'}) = \langle \Omega_\tau, \Phi_\tau(f_1) \cdots \Phi_\tau(f_n) \Phi_\tau^+(g_1) \cdots \Phi_\tau^+(g_{n'}) \Omega_\tau \rangle_\tau.$$

As in the bosonic case, the n -point functions satisfy the field equation in the distributional sense in each argument and extend to distributions on $M \times \cdots \times M$.

If one uses the self-dual fermionic algebra $\mathfrak{A}_{\text{ferm}}^{\text{sd}}(M, S, P)$ instead of $\mathfrak{A}_{\text{ferm}}(M, S, P)$, then one gets the quantum field

$$\Psi_\tau(f) := \pi_\tau(\mathbf{b}(Gf))$$

where \mathbf{b} is as in Definition A.6. Then the analogue to Theorem 4.5 is

Theorem 4.8. *Let (M, S, P) be an object in GlobHypSkewDef and τ a state on the corresponding self-dual fermionic algebra $\mathfrak{A}_{\text{ferm}}^{\text{sd}}(M, S, P)$. Then*

- (i) $P\Psi_\tau = 0$ holds in the distributional sense;
- (ii) the quantum field takes values in self-adjoint operators, $\Psi_\tau(f) = \Psi_\tau(f)^*$ for all $f \in C_c^\infty(M, S)$;
- (iii) the quantum fields satisfy the canonical anti-commutation relations, i.e.,

$$\{\Psi_\tau(f), \Psi_\tau(g)\} = \int_M \langle Gf, g \rangle dV \cdot \text{id}_{\mathcal{H}_\tau}$$

for all $f, g \in C_c^\infty(M, S)$.

Remark 4.9. It is interesting to compare the concept of locally covariant quantum field theories as proposed in [11] to the axiomatic approach to quantum field theory on Minkowski space based on the Gårding-Wightman axioms as exposed in [35, Sec. IX.8]. Property 1 (relativistic invariance of states) and Property 6 (Poincaré invariance of the field) in [35] are replaced by functoriality (covariance). Property 4 (invariant domain for fields) and Property 5 (regularity of the field) have been encoded in strong regularity of the state used to define the quantum field in the bosonic case and are automatic in the fermionic case. Property 7 (local commutativity or microscopic causality) is contained in Theorems 4.2 and 4.5. Property 3 (existence and uniqueness of the vacuum) has no analogue and is replaced by the *choice* of a state. Property 8 (cyclicity of the vacuum) is then automatic by the general properties of the GNS-construction.

There remains one axiom, Property 2 (spectral condition), which we have not discussed at all. It gets replaced by the Hadamard condition on the state chosen. It was observed by Radzikowski [32] that earlier formulations of this condition are equivalent to a condition on the wave front set of the 2-point function. Much work has been put into constructing and investigating Hadamard states for various examples of fields, see e.g. [15, 16, 19, 25, 36, 37, 38, 42] and the references therein.

APPENDIX A. ALGEBRAS OF CANONICAL (ANTI-) COMMUTATION
RELATIONS

We collect the necessary algebraic facts about CAR and CCR-algebras.

A.1. CAR algebras. The symbol ‘‘CAR’’ stands for ‘‘canonical anti-commutation relations’’. These algebras are related to pre-Hilbert spaces. We always assume the Hermitian inner product (\cdot, \cdot) to be linear in the first argument and anti-linear in the second.

Definition A.1. A *CAR-representation* of a complex pre-Hilbert space $(V, (\cdot, \cdot))$ is a pair (\mathbf{a}, A) , where A is a unital C^* -algebra and $\mathbf{a} : V \rightarrow A$ is an anti-linear map satisfying:

- (i) $A = C^*(\mathbf{a}(V))$,
- (ii) $\{\mathbf{a}(v_1), \mathbf{a}(v_2)\} = 0$ and
- (iii) $\{\mathbf{a}(v_1)^*, \mathbf{a}(v_2)\} = (v_1, v_2) \cdot 1$,

for all $v_1, v_2 \in V$.

We want to discuss CAR-representations in terms of C^* -Clifford algebras, whose definition we recall. Given a complex pre-Hilbert vector space $(V, (\cdot, \cdot))$, we denote by $V_{\mathbb{C}} := V \otimes_{\mathbb{R}} \mathbb{C}$ the complexification of V considered as a real vector space and by $q_{\mathbb{C}}$ the complex-bilinear extension of $\Re(\cdot, \cdot)$ to $V_{\mathbb{C}}$. Let $\text{Cl}_{\text{alg}}(V_{\mathbb{C}}, q_{\mathbb{C}})$ be the algebraic Clifford algebra of $(V_{\mathbb{C}}, q_{\mathbb{C}})$. It is an associative complex algebra with unit and contains $V_{\mathbb{C}}$ as a vector subspace. Its multiplication is called Clifford multiplication and denoted by ‘‘ \cdot ’’. It satisfies the Clifford relations

$$(15) \quad v \cdot w + w \cdot v = -2q_{\mathbb{C}}(v, w)1$$

for all $v, w \in V_{\mathbb{C}}$. Define the $*$ -operator on $\text{Cl}_{\text{alg}}(V_{\mathbb{C}}, q_{\mathbb{C}})$ to be the unique anti-multiplicative and anti-linear extension of the anti-linear map $V_{\mathbb{C}} \rightarrow V_{\mathbb{C}}$, $v_1 + iv_2 \mapsto -\overline{(v_1 + iv_2)} = -(v_1 - iv_2)$ for all $v_1, v_2 \in V$. In other words,

$$*\left(\sum_{i_1 < \dots < i_k} \alpha_{i_1, \dots, i_k} z_{i_1} \cdot \dots \cdot z_{i_k}\right) = (-1)^k \sum_{i_1 < \dots < i_k} \overline{\alpha_{i_1, \dots, i_k}} \cdot \overline{z_{i_k}} \cdot \dots \cdot \overline{z_{i_1}}$$

for all $k \in \mathbb{N}$ and $z_{i_1}, \dots, z_{i_k} \in V_{\mathbb{C}}$. Let $\|\cdot\|_{\infty}$ be defined by

$$\|a\|_{\infty} := \sup_{\pi \in \text{Rep}(V)} (\|\pi(a)\|)$$

for every $a \in \text{Cl}_{\text{alg}}(V_{\mathbb{C}}, q_{\mathbb{C}})$, where $\text{Rep}(V)$ denotes the set of all (isomorphism classes of) $*$ -homomorphisms from $\text{Cl}_{\text{alg}}(V_{\mathbb{C}}, q_{\mathbb{C}})$ to C^* -algebras. Then $\|\cdot\|_{\infty}$ can be shown to be a well-defined C^* -norm on $\text{Cl}_{\text{alg}}(V_{\mathbb{C}}, q_{\mathbb{C}})$, see e.g. [31, Sec. 1.2].

Definition A.2. The C^* -Clifford algebra of a pre-Hilbert space $(V, (\cdot, \cdot))$ is the C^* -completion of $\text{Cl}_{\text{alg}}(V_{\mathbb{C}}, q_{\mathbb{C}})$ with respect to the C^* -norm $\|\cdot\|_{\infty}$ and the star operator defined above.

Theorem A.3. For every complex pre-Hilbert space $(V, (\cdot, \cdot))$, the C^* -Clifford algebra $\text{Cl}(V_{\mathbb{C}}, q_{\mathbb{C}})$ provides a CAR-representation of $(V, (\cdot, \cdot))$ via $\mathbf{a}(v) = \frac{1}{2}(v + iJv)$, where J is the complex structure of V .

Moreover, CAR-representations have the following universal property: Let \widehat{A} be any unital C^* -algebra and $\widehat{\mathbf{a}} : V \rightarrow \widehat{A}$ be any anti-linear map satisfying Axioms (ii) and (iii) of Definition A.1. Then there exists a unique C^* -morphism

$\tilde{\alpha} : \text{Cl}(V_{\mathbb{C}}, q_{\mathbb{C}}) \rightarrow \widehat{A}$ such that

$$\begin{array}{ccc} V & \xrightarrow{\widehat{\mathbf{a}}} & \widehat{A} \\ \mathbf{a} \downarrow & \nearrow \tilde{\alpha} & \\ \text{Cl}(V_{\mathbb{C}}, q_{\mathbb{C}}) & & \end{array}$$

commutes. Furthermore, $\tilde{\alpha}$ is injective.

Proof. Define $p_{\mp} : V \rightarrow \text{Cl}(V_{\mathbb{C}}, q_{\mathbb{C}})$ by $p_{-}(v) := \frac{1}{2}(v + iJv)$ and $p_{+}(v) := \frac{1}{2}(v - iJv)$. Since $p_{-}(Jv) = -ip_{-}(v)$, the map $\mathbf{a} = p_{-}$ is anti-linear. Because of $\mathbf{a}(v) - \mathbf{a}(v)^* = p_{-}(v) + p_{+}(v) = v$, the \mathbb{C}^* -subalgebra of $\text{Cl}(V_{\mathbb{C}}, q_{\mathbb{C}})$ generated by the image of \mathbf{a} contains V . Hence $\mathbf{a}(V)$ generates $\text{Cl}(V_{\mathbb{C}}, q_{\mathbb{C}})$ as a \mathbb{C}^* -algebra. Axiom (i) in Definition A.1 is proved.

Let $v_1, v_2 \in V$, then

$$\begin{aligned} \{\mathbf{a}(v_1), \mathbf{a}(v_2)\} &= p_{-}(v_1) \cdot p_{-}(v_2) + p_{-}(v_2) \cdot p_{-}(v_1) \\ &= -2q_{\mathbb{C}}(p_{-}(v_1), p_{-}(v_2)) \cdot 1 \\ &= 0, \end{aligned}$$

which is Axiom (ii) in Definition A.1. Furthermore,

$$\begin{aligned} \{\mathbf{a}(v_1)^*, \mathbf{a}(v_2)\} &= -p_{+}(v_1) \cdot p_{-}(v_2) - p_{-}(v_2) \cdot p_{+}(v_1) \\ &= 2q_{\mathbb{C}}(p_{+}(v_1), p_{-}(v_2)) \cdot 1 \\ &= \Re(v_1, v_2) \cdot 1 + i\Re(v_1, Jv_2) \cdot 1 \\ &= (v_1, v_2) \cdot 1, \end{aligned}$$

which shows Axiom (iii) in Definition A.1. Therefore $(\mathbf{a}, \text{Cl}(V_{\mathbb{C}}, q_{\mathbb{C}}))$ is a CAR-representation of $(V, (\cdot, \cdot))$.

The second part of the theorem follows from $\text{Cl}(V_{\mathbb{C}}, q_{\mathbb{C}})$ being simple, i.e., from the non-existence of non-trivial closed two-sided $*$ -invariant ideals, see [31, Thm. 1.2.2]. Let $\widehat{\mathbf{a}} : V \rightarrow \widehat{A}$ be any other anti-linear map satisfying (ii) and (iii) in Definition A.1. Since \mathbf{a} and $\widehat{\mathbf{a}}$ are injective (which is clear by Axiom (iii)) one may set $\alpha(\mathbf{a}(v)) := \widehat{\mathbf{a}}(v)$ for all $v \in V$. Axioms (ii) and (iii) allow us to extend α to a \mathbb{C}^* -morphism $\tilde{\alpha} : C^*(\mathbf{a}(V)) = \text{Cl}(V_{\mathbb{C}}, q_{\mathbb{C}}) \rightarrow \widehat{A}$. The injectivity of $\widehat{\mathbf{a}}$ implies the non-triviality of $\tilde{\alpha}$ which, together with the simplicity of $\text{Cl}(V_{\mathbb{C}}, q_{\mathbb{C}})$, provides the injectivity of $\tilde{\alpha}$. Therefore we found an injective \mathbb{C}^* -morphism $\tilde{\alpha} : \text{Cl}(V_{\mathbb{C}}, q_{\mathbb{C}}) \rightarrow \widehat{A}$ with $\tilde{\alpha} \circ \mathbf{a} = \widehat{\mathbf{a}}$. It is unique since it is determined by \mathbf{a} and $\widehat{\mathbf{a}}$ on a subset of generators. This concludes the proof of Theorem A.3. \square

For an alternative description of the CAR-representation in terms of creation and annihilation operators on the fermionic Fock space we refer to [9, Prop. 5.2.2].

Corollary A.4. *For every complex pre-Hilbert space $(V, (\cdot, \cdot))$ there exists a CAR-representation of $(V, (\cdot, \cdot))$, unique up to \mathbb{C}^* -isomorphism.*

Proof. The existence has already been proved in Theorem A.3. Let $(\widehat{\mathbf{a}}, \widehat{A})$ be any CAR-representation of $(V, (\cdot, \cdot))$. Theorem A.3 states the existence of a unique injective \mathbb{C}^* -morphism $\tilde{\alpha} : \text{Cl}(V_{\mathbb{C}}, q_{\mathbb{C}}) \rightarrow \widehat{A}$ such that $\tilde{\alpha} \circ \mathbf{a} = \widehat{\mathbf{a}}$. Now $\tilde{\alpha}$ has to be surjective since Axiom (i) holds for $(\widehat{\mathbf{a}}, \widehat{A})$. \square

From now on, given a complex pre-Hilbert space $(V, (\cdot, \cdot))$, we denote the C^* -algebra $\text{Cl}(V_{\mathbb{C}}, q_{\mathbb{C}})$ associated with the CAR-representation $(\mathbf{a}, \text{Cl}(V_{\mathbb{C}}, q_{\mathbb{C}}))$ of $(V, (\cdot, \cdot))$ by $\text{CAR}(V, (\cdot, \cdot))$. We list the properties of CAR-representations which are relevant for quantization, see also [9, Vol. II, Thm. 5.2.5, p. 15].

Proposition A.5. *Let $(V, (\cdot, \cdot))$ be a complex pre-Hilbert space and $(\mathbf{a}, \text{CAR}(V, (\cdot, \cdot)))$ its CAR-representation.*

- (i) *For every $v \in V$ one has $\|\mathbf{a}(v)\| = |v| = (v, v)^{\frac{1}{2}}$, where $\|\cdot\|$ denotes the C^* -norm on $\text{CAR}(V, (\cdot, \cdot))$.*
- (ii) *The C^* -algebra $\text{CAR}(V, (\cdot, \cdot))$ is simple, i.e., it has no closed two-sided $*$ -ideals other than $\{0\}$ and the algebra itself.*
- (iii) *The algebra $\text{CAR}(V, (\cdot, \cdot))$ is \mathbb{Z}_2 -graded,*

$$\text{CAR}(V, (\cdot, \cdot)) = \text{CAR}^{\text{even}}(V, (\cdot, \cdot)) \oplus \text{CAR}^{\text{odd}}(V, (\cdot, \cdot)),$$

and $\mathbf{a}(V) \subset \text{CAR}^{\text{odd}}(V, (\cdot, \cdot))$.

- (iv) *Let $f : V \rightarrow V'$ be an isometric linear embedding, where $(V', (\cdot, \cdot)')$ is another complex pre-Hilbert space. Then there exists a unique injective C^* -morphism $\text{CAR}(f) : \text{CAR}(V, (\cdot, \cdot)) \rightarrow \text{CAR}(V', (\cdot, \cdot)')$ such that*

$$\begin{array}{ccc} V & \xrightarrow{f} & V' \\ \downarrow \mathbf{a} & & \downarrow \mathbf{a}' \\ \text{CAR}(V, (\cdot, \cdot)) & \xrightarrow{\text{CAR}(f)} & \text{CAR}(V', (\cdot, \cdot)') \end{array}$$

commutes.

Proof. We show assertion (i). On the one hand, the C^* -property of the norm $\|\cdot\|$ implies

$$\begin{aligned} \|\mathbf{a}(v)\|^4 &= \|\mathbf{a}(v)\mathbf{a}(v)^*\|^2 \\ &= \|(\mathbf{a}(v)\mathbf{a}(v)^*)^2\|. \end{aligned}$$

On the other hand,

$$\begin{aligned} (\mathbf{a}(v)\mathbf{a}(v)^*)^2 &= \mathbf{a}(v)\{\mathbf{a}(v)^*, \mathbf{a}(v)\}\mathbf{a}(v)^* \\ &= |v|^2 \mathbf{a}(v)\mathbf{a}(v)^*, \end{aligned}$$

where we used $\mathbf{a}(v)^2 = 0$ which follows from the second axiom. We deduce that

$$\begin{aligned} \|\mathbf{a}(v)\|^4 &= |v|^2 \cdot \|\mathbf{a}(v)\mathbf{a}(v)^*\| \\ &= |v|^2 \cdot \|\mathbf{a}(v)\|^2. \end{aligned}$$

Since \mathbf{a} is injective, we obtain the result.

Assertion (ii) follows from $\text{Cl}(V_{\mathbb{C}}, q_{\mathbb{C}})$ being simple, see [31, Thm. 1.2.2]. Alternatively, it can be deduced from the universal property formulated in Theorem A.3.

To see (iii) we recall that the Clifford algebra $\text{Cl}(V_{\mathbb{C}}, q_{\mathbb{C}})$ has a \mathbb{Z}_2 -grading where the even part is generated by products of an even number of vectors in $V_{\mathbb{C}}$ and, similarly, the odd part is the vector space span of products of an odd number of vectors in $V_{\mathbb{C}}$, see [31, p. 27]. This is compatible with the Clifford relations (15). Clearly, $\mathbf{a}(V) \subset \text{CAR}^{\text{odd}}(V, (\cdot, \cdot))$.

It remains to show (iv). It is straightforward to check that $\mathbf{a}' \circ f$ satisfies Axioms (ii) and (iii) in Definition A.1. The result follows from Theorem A.3. \square

One easily sees that $\text{CAR}(\text{id}) = \text{id}$ and that $\text{CAR}(f' \circ f) = \text{CAR}(f') \circ \text{CAR}(f)$ for all isometric linear embeddings $V \xrightarrow{f} V' \xrightarrow{f'} V''$. Therefore we have constructed a covariant functor

$$\text{CAR} : \text{HILB} \longrightarrow \mathbb{C}^* \text{Alg},$$

where HILB denotes the category whose objects are the complex pre-Hilbert spaces and whose morphisms are the isometric linear embeddings.

For real pre-Hilbert spaces there is the concept of *self-dual* CAR-representations.

Definition A.6. A *self-dual* CAR-representation of a real pre-Hilbert space $(V, (\cdot, \cdot))$ is a pair (\mathbf{b}, A) , where A is a unital \mathbb{C}^* -algebra and $\mathbf{b} : V \rightarrow A$ is an \mathbb{R} -linear map satisfying:

- (i) $A = \mathbb{C}^*(\mathbf{b}(V))$,
- (ii) $\mathbf{b}(v) = \mathbf{b}(v)^*$ and
- (iii) $\{\mathbf{b}(v_1), \mathbf{b}(v_2)\} = (v_1, v_2) \cdot 1$,

for all $v, v_1, v_2 \in V$.

Given a self-dual CAR-representation, one can extend \mathbf{b} to a \mathbb{C} -linear map from the complexification $V_{\mathbb{C}}$ to A . This extension $\mathbf{b} : V_{\mathbb{C}} \rightarrow A$ then satisfies $\mathbf{b}(\bar{v}) = \mathbf{b}(v)^*$ and $\{\mathbf{b}(v_1), \mathbf{b}(v_2)\} = (v_1, \bar{v}_2) \cdot 1$ for all $v, v_1, v_2 \in V_{\mathbb{C}}$. These are the axioms of a self-dual CAR-representation as in [1, p. 386].

Theorem A.7. For every real pre-Hilbert space $(V, (\cdot, \cdot))$, the \mathbb{C}^* -Clifford algebra $\text{Cl}(V_{\mathbb{C}}, q_{\mathbb{C}})$ provides a self-dual CAR-representation of $(V, (\cdot, \cdot))$ via $\mathbf{b}(v) = \frac{i}{\sqrt{2}}v$. Moreover, self-dual CAR-representations have the following universal property: Let \hat{A} be any unital \mathbb{C}^* -algebra and $\hat{\mathbf{b}} : V \rightarrow \hat{A}$ be any \mathbb{R} -linear map satisfying Axioms (ii) and (iii) of Definition A.6. Then there exists a unique \mathbb{C}^* -morphism $\tilde{\beta} : \text{Cl}(V_{\mathbb{C}}, q_{\mathbb{C}}) \rightarrow \hat{A}$ such that

$$\begin{array}{ccc} V & \xrightarrow{\hat{\mathbf{b}}} & \hat{A} \\ \mathbf{b} \downarrow & \nearrow \tilde{\beta} & \\ \text{Cl}(V_{\mathbb{C}}, q_{\mathbb{C}}) & & \end{array}$$

commutes. Furthermore, $\tilde{\beta}$ is injective.

Corollary A.8. For every real pre-Hilbert space $(V, (\cdot, \cdot))$ there exists a CAR-representation of $(V, (\cdot, \cdot))$, unique up to \mathbb{C}^* -isomorphism.

From now on, given a real pre-Hilbert space $(V, (\cdot, \cdot))$, we denote the \mathbb{C}^* -algebra $\text{Cl}(V_{\mathbb{C}}, q_{\mathbb{C}})$ associated with the self-dual CAR-representation $(\mathbf{b}, \text{Cl}(V_{\mathbb{C}}, q_{\mathbb{C}}))$ of $(V, (\cdot, \cdot))$ by $\text{CAR}_{\text{sd}}(V, (\cdot, \cdot))$.

Proposition A.9. Let $(V, (\cdot, \cdot))$ be a real pre-Hilbert space and $(\mathbf{b}, \text{CAR}_{\text{sd}}(V, (\cdot, \cdot)))$ its self-dual CAR-representation.

- (i) For every $v \in V$ one has $\|\mathbf{b}(v)\| = \frac{1}{\sqrt{2}}\|v\|$, where $\|\cdot\|$ denotes the \mathbb{C}^* -norm on $\text{CAR}_{\text{sd}}(V, (\cdot, \cdot))$.
- (ii) The \mathbb{C}^* -algebra $\text{CAR}_{\text{sd}}(V, (\cdot, \cdot))$ is simple.
- (iii) The algebra $\text{CAR}_{\text{sd}}(V, (\cdot, \cdot))$ is \mathbb{Z}_2 -graded,

$$\text{CAR}_{\text{sd}}(V, (\cdot, \cdot)) = \text{CAR}_{\text{sd}}^{\text{even}}(V, (\cdot, \cdot)) \oplus \text{CAR}_{\text{sd}}^{\text{odd}}(V, (\cdot, \cdot)),$$

and $\mathbf{b}(V) \subset \text{CAR}_{\text{sd}}^{\text{odd}}(V, (\cdot, \cdot))$.

(iv) Let $f : V \rightarrow V'$ be an isometric linear embedding, where $(V', (\cdot, \cdot)')$ is another real pre-Hilbert space. Then there exists a unique injective C^* -morphism $\text{CAR}_{\text{sd}}(f) : \text{CAR}_{\text{sd}}(V, (\cdot, \cdot)) \rightarrow \text{CAR}_{\text{sd}}(V', (\cdot, \cdot)')$ such that

$$\begin{array}{ccc} V & \xrightarrow{f} & V' \\ \downarrow \mathbf{b} & & \downarrow \mathbf{b}' \\ \text{CAR}_{\text{sd}}(V, (\cdot, \cdot)) & \xrightarrow{\text{CAR}_{\text{sd}}(f)} & \text{CAR}_{\text{sd}}(V', (\cdot, \cdot)') \end{array}$$

commutes.

The proofs are similar to the ones for CAR-representations of complex pre-Hilbert spaces. We have constructed a functor

$$\text{CAR}_{\text{sd}} : \text{HILB}_{\mathbb{R}} \longrightarrow C^* \text{Alg},$$

where $\text{HILB}_{\mathbb{R}}$ denotes the category whose objects are the real pre-Hilbert spaces and whose morphisms are the isometric linear embeddings.

Remark A.10. Let $(V, (\cdot, \cdot))$ be a complex pre-Hilbert space. If we consider V as a real vector space, then we have the real pre-Hilbert space $(V, \Re(\cdot, \cdot))$. For the corresponding CAR-representations we have

$$\text{CAR}(V, (\cdot, \cdot)) = \text{CAR}_{\text{sd}}(V, \Re(\cdot, \cdot)) = \text{Cl}(V_{\mathbb{C}}, q_{\mathbb{C}})$$

and

$$\mathbf{b}(v) = \frac{i}{\sqrt{2}}(\mathbf{a}(v) - \mathbf{a}(v)^*).$$

A.2. CCR algebras. In this section, we recall the construction of the representation of any (real) symplectic vector space by the so-called canonical commutation relations (CCR). Proofs can be found in [4, Sec. 4.2].

Definition A.11. A CCR-representation of a symplectic vector space (V, ω) is a pair (w, A) , where A is a unital C^* -algebra and w is a map $V \rightarrow A$ satisfying:

- (i) $A = C^*(w(V))$,
- (ii) $w(0) = 1$,
- (iii) $w(-\varphi) = w(\varphi)^*$,
- (iv) $w(\varphi + \psi) = e^{i\omega(\varphi, \psi)/2} w(\varphi) \cdot w(\psi)$,

for all $\varphi, \psi \in V$.

The map w is in general neither linear, nor any kind of group homomorphism, nor continuous [4, Prop. 4.2.3].

Example A.12. Given any symplectic vector space (V, ω) , consider the Hilbert space $H := L^2(V, \mathbb{C})$, where V is endowed with the counting measure. Define the map w from V into the space $\mathcal{L}(H)$ of bounded endomorphisms of H by

$$(w(\varphi)F)(\psi) := e^{i\omega(\varphi, \psi)/2} F(\varphi + \psi),$$

for all $\varphi, \psi \in V$ and $F \in H$. It is well-known that $\mathcal{L}(H)$ is a C^* -algebra with the operator norm as C^* -norm, and that the map w satisfies the Axioms (ii)-(iv) from Definition A.11, see e.g. [4, Ex. 4.2.2]. Hence setting $A := C^*(w(V))$, the pair (w, A) provides a CCR-representation of (V, ω) .

This is essentially the only example of CCR-representation:

Theorem A.13. *Let (V, ω) be a symplectic vector space and (\hat{w}, \hat{A}) be a pair satisfying the Axioms (ii)-(iv) of Definition A.11. Then there exists a unique C^* -morphism $\Phi : A \rightarrow \hat{A}$ such that $\Phi \circ w = \hat{w}$, where (w, A) is the CCR-representation from Example A.12. Moreover, Φ is injective.*

In particular, (V, ω) has a CCR-representation, unique up to C^ -isomorphism.*

We denote the C^* -algebra associated to the CCR-representation of (V, ω) from Example A.12 by $\text{CCR}(V, \omega)$. As a consequence of Theorem A.13, we obtain the following important corollary.

Corollary A.14. *Let (V, ω) be a symplectic vector space and $(w, \text{CCR}(V, \omega))$ its CCR-representation.*

- (i) *The C^* -algebra $\text{CCR}(V, \omega)$ is simple, i.e., it has no closed two-sided $*$ -ideals other than $\{0\}$ and the algebra itself.*
- (ii) *Let (V', ω') be another symplectic vector space and $f : V \rightarrow V'$ a symplectic linear map. Then there exists a unique injective C^* -morphism $\text{CCR}(f) : \text{CCR}(V, \omega) \rightarrow \text{CCR}(V', \omega')$ such that*

$$\begin{array}{ccc} V & \xrightarrow{f} & V' \\ \downarrow w & & \downarrow w' \\ \text{CCR}(V, \omega) & \xrightarrow{\text{CCR}(f)} & \text{CCR}(V', \omega') \end{array}$$

commutes.

Obviously $\text{CCR}(\text{id}) = \text{id}$ and $\text{CCR}(f' \circ f) = \text{CCR}(f') \circ \text{CCR}(f)$ for all symplectic linear maps $V \xrightarrow{f} V' \xrightarrow{f'} V''$, so that we have constructed a covariant functor

$$\text{CCR} : \text{Sympl} \longrightarrow C^* \text{ Alg.}$$

REFERENCES

- [1] H. ARAKI: *On quasifree states of CAR and Bogoliubov automorphisms.* Publ. Res. Inst. Math. Sci. **6** (1970/71), 385–442.
- [2] C. BÄR AND C. BECKER: *C^* -algebras.* In: C. Bär and K. Fredenhagen (Eds.): *Quantum field theory on curved spacetimes.* 1-37, Lecture Notes in Phys. **786**, Springer-Verlag, Berlin, 2009.
- [3] C. BÄR, P. GAUDUCHON, AND A. MOROIANU: *Generalized Cylinders in Semi-Riemannian and Spin Geometry.* Math. Zeitschr. **249** (2005), 545–580.
- [4] C. BÄR, N. GINOUX, AND F. PFÄFFLE: *Wave Equations on Lorentzian Manifolds and Quantization.* EMS, Zürich, 2007.
- [5] H. BAUM: *Spin-Strukturen und Dirac-Operatoren über pseudoriemannschen Mannigfaltigkeiten.* Teubner, Leipzig, 1981.
- [6] A. N. BERNAL AND M. SÁNCHEZ: *Smoothness of time functions and the metric splitting of globally hyperbolic spacetimes.* Commun. Math. Phys. **257** (2005), 43–50.
- [7] A. N. BERNAL AND M. SÁNCHEZ: *Further results on the smoothability of Cauchy hypersurfaces and Cauchy time functions.* Lett. Math. Phys. **77** (2006), 183–197.
- [8] T. BRANSON AND O. HIJAZI: *Bochner-Weitzenböck formulas associated with the Rarita-Schwinger operator.* Internat. J. Math. **13** (2002), 137–182.
- [9] O. BRATTELI AND D. W. ROBINSON: *Operator algebras and quantum statistical mechanics, I-II* (second edition). Texts and Monographs in Physics, Springer, Berlin, 1997.
- [10] T. BRÖCKER AND T. TOM DIECK: *Representations of compact Lie groups.* Graduate Texts in Mathematics **98**, Springer-Verlag, New York, 1995.
- [11] R. BRUNETTI, K. FREDENHAGEN AND R. VERCH: *The generally covariant locality principle - a new paradigm for local quantum field theory.* Commun. Math. Phys. **237** (2003), 31–68.

- [12] H. A. BUCHDAHL: *On the compatibility of relativistic wave equations in Riemann spaces. II.* J. Phys. A **15** (1982), 1-5.
- [13] H. A. BUCHDAHL: *On the compatibility of relativistic wave equations in Riemann spaces. III.* J. Phys. A **15** (1982), 1057-1062.
- [14] C. DAPPIAGGI, T.-P. HACK AND N. PINAMONTI: *The extended algebra of observables for Dirac fields and the trace anomaly of their stress-energy tensors.* Rev. Math. Phys. **21** (2009), 1241–1312.
- [15] C. DAPPIAGGI, V. MORETTI AND N. PINAMONTI: *Distinguished quantum states in a class of cosmological spacetimes and their Hadamard property.* J. Math. Phys. **50** (2009), 062304, 38 p.
- [16] C. DAPPIAGGI, N. PINAMONTI AND M. PORRMANN: *Local causal structures, Hadamard states and the principle of local covariance in quantum field theory.* arXiv:1001.0858
- [17] J. DIMOCK: *Algebras of local observables on a manifold.* Commun. Math. Phys. **77** (1980), 219–228.
- [18] J. DIMOCK: *Dirac quantum fields on a manifold.* Trans. Amer. Math. Soc. **269** (1982), 133–147.
- [19] C. J. FEWSTER AND R. VERCH: *A quantum weak energy inequality for Dirac fields in curved spacetime.* Commun. Math. Phys. **225** (2002), 331–359.
- [20] E. FURLANI: *Quantization of massive vector fields in curved space-time.* J. Math. Phys. **40** (1999), 2611-2626.
- [21] R.P. GEROGH: *Domain of dependence.* J. Math. Phys. **11** (1970), 437–449.
- [22] G. W. GIBBONS: *A note on the Rarita-Schwinger equation in a gravitational background.* J. Phys. A **9** (1976), 145–148.
- [23] L. HÖRMANDER: *The analysis of linear partial differential operators. I. Distribution theory and Fourier analysis.* 2nd ed. Grundlehren der Mathematischen Wissenschaften **256**, Springer-Verlag, Berlin, 1990.
- [24] L. HÖRMANDER: *The analysis of linear partial differential operators. III. Pseudodifferential operators.* Grundlehren der Mathematischen Wissenschaften **274**, Springer-Verlag, Berlin, 1985.
- [25] S. HOLLANDS AND R. M. WALD: *Axiomatic quantum field theory in curved spacetime.* Commun. Math. Phys. **293** (2010), 85–125.
- [26] B. S. KAY: *Linear spin-zero quantum fields in external gravitational and scalar fields.* Commun. Math. Phys. **62** (1978), 55–70.
- [27] H. B. LAWSON AND M.-L. MICHELSON: *Spin Geometry.* Princeton University Press, Princeton, 1989.
- [28] R. MÜHLHOFF: *Higher Spin fields on curved spacetimes.* Diplomarbeit, Universität Leipzig, 2007.
- [29] R. MÜHLHOFF: *Cauchy Problem and Green's Functions for First Order Differential Operators and Algebraic Quantization.* J. Math. Phys. **52** (2011), 022303, 7 p.
- [30] B. O'NEILL: *Semi-Riemannian Geometry.* Academic Press, San Diego, 1983.
- [31] R. J. PLYMEN AND P.L. ROBINSON: *Spinors in Hilbert space.* Cambridge Tracts in Mathematics **114**, Cambridge University Press, Cambridge, 1994.
- [32] M. J. RADZIKOWSKI: *Micro-local approach to the Hadamard condition in quantum field theory on curved space-time.* Commun. Math. Phys. **179** (1996), 529-553.
- [33] W. RARITA AND J. SCHWINGER: *On a Theory of Particles with Half-Integral Spin.* Phys. Rev. **60** (1941), 61.
- [34] M. REED AND B. SIMON: *Methods of Modern Mathematical Physics I: Functional Analysis.* Academic Press, Orlando, 1980.
- [35] M. REED AND B. SIMON: *Methods of Modern Mathematical Physics II: Fourier Analysis, Self-Adjointness.* Academic Press, Orlando, 1975.
- [36] H. SAHLMANN AND R. VERCH: *Passivity and microlocal spectrum condition.* Commun. Math. Phys. **214** (2000), 705–731.
- [37] H. SAHLMANN AND R. VERCH: *Microlocal spectrum condition and Hadamard form for vector-valued quantum fields in curved spacetime.* Rev. Math. Phys. **13** (2001), 1203–1246.
- [38] K. SANDERS: *The locally covariant Dirac field.* Rev. Math. Phys. **22** (2010), 381–430.
- [39] A. STROHMAIER: *The Reeh-Schlieder property for quantum fields on stationary spacetimes.* Commun. Math. Phys. **215** (2000), 105–118.

- [40] M. E. TAYLOR: *Partial Differential Equations I - Basic Theory*. Springer-Verlag, New York - Berlin - Heidelberg, 1996.
- [41] R. VERCH: *A spin-statistics theorem for quantum fields on curved spacetime manifolds in a generally covariant framework*. *Commun. Math. Phys.* **223** (2001), 261–288.
- [42] R. M. WALD: *Quantum field theory in curved spacetime and black hole thermodynamics*. University of Chicago Press, Chicago, 1994.
- [43] MCK. Y. WANG: *Preserving parallel spinors under metric deformations*. *Indiana Univ. Math. J.* **40** (1991), 815–844.
- [44] V. WÜNSCH: *Cauchy's problem and Huygens' principle for relativistic higher spin wave equations in an arbitrary curved space-time*. *Gen. Relativity Gravitation* **17** (1985), 15-38.

UNIVERSITÄT POTSDAM, INSTITUT FÜR MATHEMATIK, AM NEUEN PALAIS 10, HAUS 8, 14469 POTSDAM, GERMANY

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT REGENSBURG, 93040 REGENSBURG, GERMANY
E-mail address: baer@math.uni-potsdam.de
E-mail address: nicolas.ginoux@mathematik.uni-regensburg.de