

Julia Pawlak | Dominique Christian Noetzel | Claudia Drago | Guntram Weithoff

# Assessing the toxicity of polystyrene beads and silica particles on the microconsumer *Brachionus calyciflorus* at different timescales

**Suggested citation referring to the original publication:**

Frontiers in Environmental Science (2022), pp. 1 - 11

DOI <https://doi.org/10.3389/fenvs.2022.955425>

ISSN 2296-665X

**Journal article | Version of record**

Secondary publication archived on the Publication Server of the University of Potsdam:

Zweitveröffentlichungen der Universität Potsdam :

Mathematisch-Naturwissenschaftliche Reihe 1277

ISSN: 1866-8372

<https://nbn-resolving.org/urn:nbn:de:kobv:517-opus4-569967>

DOI: <https://doi.org/10.25932/publishup-56996>

**Terms of use:**

This work is licensed under a Creative Commons License. This does not apply to quoted content from other authors. To view a copy of this license visit

<https://creativecommons.org/licenses/by/4.0/>.





## OPEN ACCESS

## EDITED BY

Anita Mukherjee,  
University of Calcutta, India

## REVIEWED BY

S.S.S. Sarma,  
Universidad Nacional Autónoma de  
México, Mexico  
Maranda Esterhuizen,  
University of Helsinki, Finland

## \*CORRESPONDENCE

Julia Pawlak,  
julia.pawlak.1@uni-potsdam.de

<sup>†</sup>These authors have contributed equally  
to this work and share first authorship

## SPECIALTY SECTION

This article was submitted to  
Toxicology, Pollution and the  
Environment,  
a section of the journal  
Frontiers in Environmental Science

RECEIVED 28 May 2022

ACCEPTED 28 June 2022

PUBLISHED 09 August 2022

## CITATION

Pawlak J, Noetzel DC, Drago C and  
Weithoff G (2022), Assessing the toxicity  
of polystyrene beads and silica particles  
on the microconsumer *Brachionus*  
*calyciflorus* at different timescales.  
*Front. Environ. Sci.* 10:955425.  
doi: 10.3389/fenvs.2022.955425

## COPYRIGHT

© 2022 Pawlak, Noetzel, Drago and  
Weithoff. This is an open-access article  
distributed under the terms of the  
[Creative Commons Attribution License  
\(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or  
reproduction in other forums is  
permitted, provided the original  
author(s) and the copyright owner(s) are  
credited and that the original  
publication in this journal is cited, in  
accordance with accepted academic  
practice. No use, distribution or  
reproduction is permitted which does  
not comply with these terms.

# Assessing the toxicity of polystyrene beads and silica particles on the microconsumer *Brachionus calyciflorus* at different timescales

Julia Pawlak<sup>1\*†</sup>, Dominique C. Noetzel<sup>1†</sup>, Claudia Drago<sup>1</sup> and  
Guntram Weithoff<sup>1,2</sup>

<sup>1</sup>Department for Ecology and Ecosystem Modelling, Institute of Biochemistry and Biology, University of Potsdam, Potsdam, Germany, <sup>2</sup>Berlin-Brandenburg Institute of Advanced Biodiversity Research (BBIB), Berlin, Germany

Environmental pollution by microplastics has become a severe problem in terrestrial and aquatic ecosystems and, according to actual prognoses, problems will further increase in the future. Therefore, assessing and quantifying the risk for the biota is crucial. Standardized short-term toxicological procedures as well as methods quantifying potential toxic effects over the whole life span of an animal are required. We studied the effect of the microplastic polystyrene on the survival and reproduction of a common freshwater invertebrate, the rotifer *Brachionus calyciflorus*, at different timescales. We used pristine polystyrene spheres of 1, 3, and 6  $\mu\text{m}$  diameter and fed them to the animals together with food algae in different ratios ranging from 0 to 50% nonfood particles. As a particle control, we used silica to distinguish between a pure particle effect and a plastic effect. After 24 h, no toxic effect was found, neither with polystyrene nor with silica. After 96 h, a toxic effect was detectable for both particle types. The size of the particles played a negligible role. Studying the long-term effect by using life table experiments, we found a reduced reproduction when the animals were fed with 3  $\mu\text{m}$  spheres together with similar-sized food algae. We conclude that the fitness reduction is mainly driven by the dilution of food by the nonfood particles rather than by a direct toxic effect.

## KEYWORDS

microplastics, rotifer, freshwater, natural particle, toxicity, environmental pollution

## 1 Introduction

Scientific and public interest in microplastics (< 5 mm) highly increased in recent years and with it the upcoming concerns of the negative impacts on wildlife and whole ecosystems. Since surveys in marine ecosystems started to evaluate plastic occurrences, first accidentally (Carpenter and Smith, 1972) and later intentionally (Eriksen et al., 2014; Kanhai et al., 2017), alarming results have been reported, and the consequences of mismanaged plastic waste dumping and the associated releases of plastic into the environment have become apparent (Fok et al., 2019). However, (micro)plastics are not only found in aquatic ecosystems. They seem to be ubiquitous; whether in the sediments of fields (Piehl et al., 2018), in the air (Allen et al., 2019), or even in the snow of the Arctic (Bergmann et al., 2019), microplastics (MP) can be found. The associated risk of this pollutant is still unknown.

Many studies reported that a wide range of marine and freshwater organisms are able to ingest MP (Naidu et al., 2018; Windsor et al., 2019). Effects of MP exposure and ingestion can be manifold and likely depend on plastic type (Zimmermann et al., 2020; Phothakwanpracha et al., 2021), particle shape (Frydkjær et al., 2017; Qiao et al., 2019), particle size, and MP concentration (Jeong et al., 2016; Sun et al., 2019). In addition, the feeding strategy (Scherer et al., 2017) and the duration of exposure (Eltemsah and Böhn, 2019; Cappello et al., 2021) are also determinants of the effect. Results of these laboratory studies are inconclusive, and a general assessment about the effect of MP on animal fitness or the risk of physiological impairments cannot be made yet. Studies reported adverse effects after MP exposure, resulting in increased mortality (Maes et al., 2020; An et al., 2021), decreased reproductive performance (Jeong et al., 2016; Jaikumar et al., 2019), decreased food intake (Cole et al., 2015; An et al., 2021), or even impacts at the molecular level (Gambardella et al., 2017). However, other studies found no or negligible effects after MP exposure on life history traits (Imhof et al., 2017; Wang et al., 2019). These inconsistent MP exposure results may be due to various reasons, such as exposure times, particle types, or test organism. For example, the duration of MP exposure differentially affects life history traits as shown in the study by Eltemsah and Böhn (2019) with the cladoceran *Daphnia magna*. While in the short-term experiment, no acute toxic effects could be observed after 48 h, the long-term experiment (up to 77 days) revealed reduced growth and shifts in the timing of reproduction (Eltemsah and Böhn, 2019). In the study by Cole et al. (2015) with the marine copepod *Calanus helgolandicus*, it is also evident that the duration of MP exposure plays an important role. Here, increased mortality, reduced egg size, and decreased hatching success were only seen later in the course of the experiment (Cole et al., 2015).

In addition to this unclear picture of the potential effects of MP on aquatic organisms, a shortcoming of current MP

research is the frequent lack of an adequate control of inorganic particles (Ogonowski et al., 2018). Aquatic organisms are exposed to a wide range of natural particles, such as particulate organic matter (Wilkinson et al., 2013) or inorganic matter, such as calcite (Stabel, 1986) or clay particles (Tietjen et al., 2005). Therefore, for MP risk assessment distinguishing between changes in life history traits caused by a specific plastic effect or a general particle effect is crucial (Connors et al., 2017). As shown in some studies, increasing concentrations of inorganic particles impair algal uptake by cladocerans (Arruda et al., 1983; Hart, 1988) and result in lower population growth (Kirk, 1991). Furthermore, inorganic particles affect zooplankton groups differentially, as shown in a competition experiment with cladocerans and rotifers by Kirk and Gilbert (1990). There, rotifers were found to be more competitive than cladocerans at high inputs of inorganic particles, likely due to their differing feeding modes (Kirk and Gilbert, 1990). Inorganic particles have rarely been used as reference particles in experiments with cladoceran (Schür et al., 2020; Zimmermann et al., 2020), and for rotifers, only one study recently included inorganic particles in MP experiments in its experimental design (Drago and Weithoff, 2021).

Nevertheless, the uptake of MP at the beginning of the food chain is considered particularly critical due to the risk of bioaccumulation and biomagnification between trophic levels (Setälä et al., 2014). Although the focus changed somewhat in recent years, most MP studies focused on marine environments, so freshwater systems are still underrepresented (Wagner et al., 2014; Blettler et al., 2018). The magnitude of MP concentrations in freshwater ecosystems has been described in several studies for lakes (Fischer et al., 2016; Di and Wang, 2018) and rivers (Mani et al., 2015; Kapp and Yeatman, 2018), demonstrating the need to study the impact of MP on aquatic freshwater organisms. In freshwater systems, rotifers are a pivotal component of zooplankton communities (Arndt, 1993), providing an important link between primary producers and secondary consumers (Walz, 1993). Also, if food resources are adequate, they are widespread and can develop high population densities in freshwaters (Wallace and Snell, 2010). Although rotifers play an important role in limnetic food webs, studies related to the impact of MP to this phylum are still extremely scarce.

The aim of this study was to investigate the effect of polystyrene MP on the life history traits reproduction and survival of the rotifer *Brachionus calyciflorus*. Therefore, we tested three MP sizes (1, 3, 6 µm) of spherical shape and exposed the rotifer, first, in a short-term acute toxicity test by using increasing MP concentrations and food limitation as additional stressor. In this approach, we used silica beads of the same size range as MP as inorganic reference particles. Second, we tested the impact of the same MP spheres in a long-term chronic toxicity test by using one MP

concentration, no reference particles, and food saturation. Thus, we excluded additional stress by food limitation to attribute any effect on life parameters to MP exposure. We chose these two types of particles because polystyrene is one of the most common plastic types found in aquatic ecosystems (Andrady, 2011; Mani et al., 2015), and silica is one of the most abundant minerals of the Earth's crust and is formed during the weathering of rock formations.

## 2 Materials and methods

### 2.1 Test particles

To examine the impact of microplastics and more natural particles on life history traits of an aquatic microconsumer, we used polystyrene (PS) and silica (Si) microsized beads. PS was purchased from Polysciences Europe GmbH, green fluorescently (excitation 441 nm, emission 486 nm) with a density of  $1.05 \text{ g cm}^{-3}$ . Si was purchased from microParticles GmbH Berlin, nonfluorescent with a density of  $1.85 \text{ g cm}^{-3}$ . We used the Si particle as a reference particle for PS only in the short-term exposure experiments. Both test particles demonstrated diameters of approximately 1.0, 3.0, and  $6.0 \mu\text{m}$  (detailed information in supplementary material, [Supplementary Table S1](#)), respectively. The particle stocks were prepared in MilliQ water in the required concentrations under sterile conditions and were stored in a fridge at  $6^\circ\text{C}$  in the dark to prevent bacterial growth. Particle concentrations were microscopically quantified using a hemocytometer (Neubauer improved; CH 30, Olympus Europa SE & Co. KG) and an Olympus (Europa SE & Co. KG) microscope. To avoid particle aggregation, the stocks were sonicated (Sonorex Digitec DT 514 BH, Bandelin electronic GmbH & Co. KG) 30 min before use. For the short-term acute toxicity test (AT, see [section 2.4.1](#)), we used the surfactant Tween 20 to prevent the aggregation of particles during the exposure time. Therefore, we added  $10 \mu\text{L}$  of 1% surfactant Tween20-stock to the particle stocks.

### 2.2 Cultivation of rotifer and algae

For the experiments, we used the parthenogenetic monogonont rotifer *B. calyciflorus* s.s. (Strain USA, [Paraskevopoulou et al., 2018](#)) as a microconsumer, which was originally collected at lake Michigan and has been in continuous culture since 2001 in our lab. Rotifer cultures for the short-term AT were fed with the spherical, unicellular green alga *Chlorella vulgaris* (Chlorophyta; strain 211-11b, Culture Collection of Algae, University of Göttingen, Germany, mean equivalent spherical diameter =  $3.4\text{--}3.7 \mu\text{m}$ ). To exclude interference of the particle shape between nonfood and food particle in the AT, we decided to use the spherical alga *C. vulgaris*. For the long-term

chronical toxicity test (CT), rotifers were fed with the sickle-shaped chlorophyte alga *Monoraphidium minutum* (strain 243-1, Culture Collection of Algae, University of Göttingen, Germany, mean equivalent spherical diameter =  $3.4 \mu\text{m}$ ). Both algae are in the same size range, of similar nutritional quality for *B. calyciflorus*, and are frequently used in laboratory experiments (*C. vulgaris*: [Sarma et al., 2000](#), *M. minutum*: [Schälicke et al., 2019](#)). Rotifers were fed *ad libitum* every week with the respective algae. Algae and rotifers were kept in modified Woods Hole (WC) Medium ([Guillard and Lorenzen, 1972](#)) at  $20^\circ\text{C}$ , at a day/night cycle of 16:8 h, and at light intensity of  $35 \mu\text{mol m}^{-2} \text{ s}^{-1}$  photosynthetic active radiation. Algal cultures were grown in semi continuous batch cultures, to keep them in exponential growth phase. Rotifer cultures were regularly fed and transferred into fresh flasks.

### 2.3 Food concentrations at acute and chronic exposure

Next, we used different food levels for acute and chronic toxicity experiments. For the AT tests, we aimed for a food level that allows for moderate growth. No food would lead to 100% mortality after 4 days making it impossible to estimate toxicity effects of MP in relation to the control. Saturating (high) food concentrations might lead to aggregations interfering with the MP. Therefore, the food concentration at which population growth is half maximal was determined by measuring population growth at cell concentrations of ( $0.05 \times 10^6$ ,  $0.125 \times 10^6$ ,  $0.25 \times 10^6$ ,  $0.5 \times 10^6$ ,  $1 \times 10^6$ ,  $2 \times 10^6 \text{ cells mL}^{-1}$ ) for 4 days. After fitting a saturation curve to the growth rate vs. food concentration data, the food concentration of half-maximal growth of  $0.41 \text{ days}^{-1}$  was  $1.43 \times 10^5 \text{ cells mL}^{-1}$  ( $\sim 2.94 \times 10^6 \mu\text{m}^3 \text{ mL}^{-1}$ ). For the CT, we wanted to test for a sublethal effect in relation to optimal environmental conditions. Thus, nonlimiting food conditions were chosen by using an algal concentration of  $5.0 \times 10^5 \text{ algae cells mL}^{-1}$  ( $\sim 14.1 \times 10^6 \mu\text{m}^3 \text{ mL}^{-1}$ ) of *M. minutum* ([Fussmann et al., 2005](#)). While MP studies often use mass or particle number as unit, to better compare treatments with the silica beads, we decided to use (bio) volume, because of their high specific weight. Previous studies demonstrated that *B. calyciflorus* ingests both nonfood particles ([Drago et al., 2020](#); [Drago and Weithoff, 2021](#)); however, we did not quantify the actual ingestion in the present study.

### 2.4 Short-and long-term test approaches

We performed two short-term acute toxicity tests, one for 24 h (AT<sub>24</sub>) and one for 96 h (AT<sub>96</sub>), and one life table experiment (CT) to assess the whole life cycle. For the AT<sub>24/96</sub> tests, we tested for the effect of three sizes of PS and silica in comparison to a control without artificial particles (only algae).

For the life table experiments, we tested only PS, because the experimental set-up did not allow for sufficient turbulence to keep the silica beads in suspension as the PS did (see below).

#### 2.4.1 Acute toxicity test (24 and 96 h)

In order to start the experiment with young animals of the same age, 4 days prior to the experiment, a subsample from the stock culture was transferred in six-well plates, each well containing 2 ml WC-Medium. Two days later, the animals were fed *ad libitum* with *C. vulgaris* to synchronize parthenogenetic reproduction. At experimental day 0, to remove remaining algae, adult rotifers were sieved through a 30  $\mu\text{m}$  mesh (Macs SmartStrainer). Thereafter, subitaneous eggs were separated from the adult rotifers by using a vortexer (Vortex Genie 2) at maximum rotation velocity for approximately 1 min. Unattached subitaneous eggs in progressive state of development (Castellanos Paez et al., 1988) were selected to gain neonates of approximately same age for the experiment (~2 h hatching span). To examine the short-term effects, *B. calyciflorus* was exposed to varying MP/silica particle concentrations. We applied a substitutive design by keeping the total volume of particles constant and varied the relative contribution of artificial food particle to the total amount of particles. The applied artificial particle concentrations were 0% (control, *C. vulgaris*:  $2.94 \times 10^6 \mu\text{m}^3 \text{mL}^{-1}$ ), 20%, 33%, 43%, and 50% proportion of artificial particle to total particles, see supplementary material, [Supplementary Table S2](#). Next, the AT<sub>24</sub> test was performed with three and the AT<sub>96</sub> test with four replicates. To prevent the aggregation of particles during the exposure time, we added 10  $\mu\text{L}$  of 1% surfactant Tween20-stock to the particle stocks. The same amount of Tween was additionally added to MilliQ water stocks. To obtain the same concentration of Tween in each glass vials, these stocks were used for the lower particle volumes. To exclude hidden effects from the surfactant agent on *B. calyciflorus*, a control with Tween and food algal for both short-term tests were made. Since we found no effect of Tween, we pooled the data from the control and the Tween treatment. This ends up in a total of 78 vials for the 24-h and 104 vials for the 96-h test.

Soda-lime glass vials (Height 24 mm, Diameter 24 mm, Volume 7 ml, Carl Roth) were filled with WC-Medium (5 ml), the appropriate particle concentration, and the food alga. Each vial was shaken for 5 s (Vortex Genie 2), and then, detached subitaneous eggs were added to the vials. To prevent air bubbles, the remaining volume was filled with WC-Medium. Subsequently, glass vials were darkened with duct tape and fixed randomly on a plankton wheel for 24 h or 96 h at a rotational speed of 13 rpm at 20°C ( $\pm 1^\circ\text{C}$ ). The vials were darkened to prevent algal growth during the exposure time. To determine the survival and growth rate of the rotifers after the end of each experiment, the content of each vial was transferred in a well of a six-well plate and rotifers (live and

dead), and their eggs were counted using a Stereomicroscope (Stemi 2000 C Carl Zeiss). Animals were declared dead if no cilia nor mastax movement was observed for 5 s. The intrinsic population growth rate  $r$  (per day) was calculated according to the equation:


$$r = \frac{\ln(N_1 + 1) - \ln(N_0)}{\Delta t} \quad (1)$$

where  $N_0$  describes the number of rotifers at the beginning of the experiment,  $N_1$  the number of rotifers at the end of the experiment, and  $\Delta t$  the time interval (4 days). Since in about one-third of the samples, no live rotifers were present after 96 h, we used the  $N_1 + 1$  transformation. No growth was detected after 24 h.

#### 2.4.2 Chronical toxicity test: Life table experiment

Three days before the start of the CT, to ensure equal and good physiological conditions, the rotifers were fed daily with approximately  $1 \times 10^6$  cells  $\text{mL}^{-1}$  of the food algal *M. minutum*. From this stock, 24 egg-bearing females were transferred into 24-well microtiter plates with the target experimental condition. The experiment started with the hatching of the neonates in their respective experimental condition. In all experiments (control and treatments), animals were provided with algal food at a concentration of  $5 \times 10^5$  cells  $\text{mL}^{-1}$  ( $\sim 14.1 \times 10^6 \mu\text{m}^3 \text{mL}^{-1}$ ). For the PS treatment, a fixed particle volume of  $9.52 \times 10^5 \mu\text{m}^3 \text{mL}^{-1}$  ( $\sim 1 \mu\text{g mL}^{-1}$ ) was used. Thus, the number of beads decreased with increasing bead sizes. The total volume of the algae was ca. 15 times higher than the volume of the beads ([Supplementary Table S2](#)).

Every 24 h ( $\pm 1$  h), the surviving rotifers were transferred in to a new well with the target conditions, and the number of eggs and neonates was recorded. The experiments were terminated when the initial rotifer died. Rotifers were declared dead when neither cilia nor mastax movement was detected after 5 s. Next, the experiment was run at 20°C ( $\pm 1^\circ\text{C}$ ) in the dark to prevent algal growth, and the plates were placed randomly on a rocker shaker to keep PS in suspension. The experiments were run sequentially, and the plastic-free control was repeated in each run. Since no differences among the different controls were found, these data were pooled. The life table experiments provided data on life span and reproduction: the average life span, the age-specific survival ( $l_x$ ), and the age-specific fecundity ( $m_x$ ) were calculated;  $x$  was defined as the age level per day,  $l_x$  as the proportion of surviving animals at the beginning of an age level, and  $m_x$  as the number of offspring per surviving animal (Begon et al., 1996). The age-specific net fecundity ( $l_x m_x$ ), the net reproduction rate ( $R_0$ , Eq. (2)), the generation time ( $T$ , Eq. 3), and the intrinsic population growth rate  $r$  (Eq. 1) were calculated as well.



$$R_0 = \sum_0^{\infty} l_x m_x \quad (2)$$

$$T = \frac{\sum x l_x m_x}{\ln(R_0)} \quad (3)$$

The intrinsic population growth rate  $r$  was calculated using the bootstrapping with 199 iterations (Weithoff and Wacker, 2007). To calculate the standard error, the 95% confidence interval obtained by the bootstrapping method was used (Higgins et al., 2019). For better comparability between results from the AT<sub>96</sub> and CT,  $r$  was calculated after the first 96 h. To prevent zeros, the data were  $N_i + 1$  transformed.

## 2.5 Statistical analyses


All data (AT and CT) were tested for normal distribution and homoscedasticity. We used One-way ANOVA for data analysis. If test requirements did not apply, nonparametric Kruskal–Wallis (KW) test or the Scheirer–Ray–Hare (SRH) test (Scheirer et al., 1976) was conducted. When significant results ( $p \leq 0.05$ ) of the above tests were obtained, Tukey HSD and Dunn’s test with Bonferroni correction were used as posthoc procedures. In general, we tested the influence of nonfood particles, size of the particles, the concentrations, and the time (only AT)

as well as the interaction of these parameters on the life history traits of *B. calyciflorus*. In the AT, we tested specifically, if differences between both nonfood-particle types could be found.

## 3 Results

### 3.1 Survival

Survival ( $N_{24}$ ) of *B. calyciflorus* after 24 h (AT<sub>24</sub>) was neither affected by PS nor by Si particles of any size or any tested particle density (Figure 1A, SRH test:  $p > 0.05$ , KW test:  $p > 0.1$ , Supplementary Table S3, S4). After 96 h (AT<sub>96</sub>), we also found no effect of particle type (SRH test:  $p = 0.9$ ) or particle size (SRH test:  $p = 0.73$ ) on rotifer survival (Supplementary Table S5). However, a variable decrease in survival was observed in response to artificial particles compared to the control. In detail, exposure to nonfood particles (PS, Si) induced a decrease in survival  $N_{96}$  of the rotifers (Figure 1B) for one size of PS and for all three sizes of Si compared to the control group (KW test: PS:  $p = 6 \mu\text{m} = 0.01$ ; Si:  $p = 1, 3, 6 \mu\text{m} = 0.02$ , Supplementary Table S6). However, no clear trend emerged with respect to the increase of nonfood proportions (Figure 1B). While no significant differences from the control were found for 1 and 3 μm PS




with increasing concentrations (KW test:  $p = 1 \mu\text{m} = 0.1$ ,  $3 \mu\text{m} = 0.08$ ), a decrease in rotifers was observed with Si particles for these two sizes: at 50% and 20% food/nonfood ratio (Dunn's test:  $p = 1 \mu\text{m} = 0.01$ ,  $3 \mu\text{m} = 0.03$ , respectively). Interestingly, survival was significantly different from control for the 6  $\mu\text{m}$  for both particle types at the lowest (20%) (Dunn's test:  $P = \text{PS} = 0.03$ ,  $\text{Si} = 0.04$ ) and highest food/nonfood proportion (50%) (Dunn's test:  $P = \text{PS}/\text{Si} = 0.02$ , [Supplementary Table S7](#)).

In CT, instead, no significant difference in survival of *B. calyciflorus* could be detected in the PS treatments compared to the control ([Figures 2A,C](#)). The average life span varied nonsignificantly between 5.0 and 6.6 days among treatments. (One-way ANOVA: for PS presence,  $p = 0.0957$ , [Supplementary Table S8](#)).

### 3.2 Reproduction and growth rate

No reproduction occurred in the AT<sub>24</sub>; thus, the intrinsic rate of population growth  $r$  was calculated only for AT<sub>96</sub>, and CT<sub>96</sub> ( $r$  of the CT was calculated after 96 h for reasons of comparison). While the growth rate for the control of the AT<sub>96</sub> was  $0.22 (\pm 0.06\text{SE}) \text{ day}^{-1}$ , the mean growth rate in both particle treatments (AT<sub>96</sub>) was either negative, or slightly above zero, for the 3 and 6  $\mu\text{m}$  Si treatments ([Supplementary Table S13](#)). No effect of the particle sizes (SRH test:  $p = 0.72$ ), the particle type (SRH test:  $p = 0.89$ ), the interaction of the two (SRH test:  $p = 0.57$ , [Supplementary Table S9](#)) was found for either nor for increasing concentrations (SRH test:  $p = \text{PS} = 0.32$ ,  $\text{Si} = 0.12$ , [Supplementary Table S10](#)). We only found effects on the intrinsic population growth rate  $r$  compared to the control





**FIGURE 3**

Intrinsic rate of population growth  $r$  (day<sup>-1</sup>) of *B. calyciflorus* after exposure to polystyrene and silica particles of three sizes (1, 3, and 6 μm) and different concentrations (0–50% food/nonfood proportion) and the control (C, food-particle only) for 96 h (AT<sub>96</sub>). Data were expressed as mean ± standard error.

(Figure 3, Supplementary Table S11). For the AT<sub>96</sub> and the CT test, different food levels were tested, and thus, the effects of PS are compared in the light of presence and absence of food limitation (Figure 4). The food levels resulted in a growth rate of 0.71 (±0.07 SE) at high food and of 0.22 (±0.06 SE) at low food, which is comparable to other studies using *B. calyciflorus* and these two food algae (Rothaupt, 1990; Weithoff and Walz, 1995; Sarma et al., 2001). In general, growth rates in CT at high food were significant higher as in the AT<sub>96</sub> at low food (KW test:  $p = 0.02$ , Supplementary Table S12). The age-specific net fecundity (Figure 2B) showed a delayed reproduction for 1 μm PS and a decreased total reproduction for 3 μm PS compared to the control. Averaged egg production (mean eggs per female: 1 μm = 9.6, 3 μm = 6.2, 6 μm = 9.0, control = 10.6) in the CT differed significantly between 3 μm and the control group (TukeyHSD:  $p = 0.001$ , Figure 2D, Supplementary Table S8). Detailed results of the CT test are listed in supplementary material (Supplementary Table S14).

## 4 Discussion


### 4.1 No acute toxicity of polystyrene

Our results indicate that short-term exposure (AT<sub>24</sub>) to PS in the size classes 1, 3, and 6 μm and at varying concentrations (6, 20, 33, 43 and 50% nonfood-particle proportion) did not demonstrate an acute toxic effect on the rotifer *B. calyciflorus*. These findings are confirmed by other studies for PS and for a range of aquatic organisms (Gambardella et al., 2018; Eltemsah and Böhn, 2019). The assumption exists that microplastic, especially of spherical shape, can be easily ingested but also egested (Cole et al., 2013; Scherer et al., 2017). Nevertheless, we

found an impact of microplastics on *B. calyciflorus* at longer timescales. The reproduction decreased in comparison with the control groups both, in the AT<sub>96</sub> and the CT test, whereas the survival was only affected in the AT<sub>96</sub> test. As reported in other studies, it seems that the reproductive output of aquatic invertebrates is more affected by the uptake of microplastic than their survival (Lee et al., 2013; Jaikumar et al., 2019). The pattern of reduced reproduction along with no clear effect on life span can be explained ecophysiologicaly. Many animals including rotifers (Weithoff, 2007; Snell, 2014) respond to a reduced food supply with a prolongation in life span on the expense of reproduction. A shortening in life span will only occur at exceptionally severe food limitation close to starvation. Since many studies do not cover the whole life cycle, changes in life span (in particular extensions) cannot be detected. In our study, a tendency from this ecophysiological pattern can be seen in the 1 μm treatment. In the first days, reproduction was slightly delayed, and survival was still relatively high, while at the peak of reproduction, it then continuously decreased.

### 4.2 Particle vs. microplastic effect

Between PS and Si, no significant effects were found neither with increasing particle size nor with increasing concentration. The results of the AT<sub>96</sub> suggest a particle rather than a microplastic effect. Similar life history responses were determined in a study by Ogonowski et al. (2016) with *D. magna* when comparing pristine PE beads with kaolin. Also, Yap et al. (2020), who used red clay as natural particles and PVC to study a potential effect of byssus production, respiration, and survival rate in a marine mussel, found no difference between the two particle types. Next, interestingly, Coady et al. (2020) found, in their study, with *D. magna* no differences at all in the investigated life history traits between



control (laboratory dilution water), the reference particles (fumed nanosized silica), and the used microplastic (nanosized ethylene acryl acid copolymer). Next, contrary, Harris and Carrington (2019) and Zimmermann et al. (2020) reported, in their studies, with the mussel *Mytilus trossulus* and the cladoceran *D. magna* that MP, but not the tested natural particles, negatively affected the life parameters of these organisms. The reasons for the different results might be manifold, for example due to the type, size, or shape of the particle tested, the concentration used, the organisms, the duration of the experiment, the observed life history traits, or the experimental design. Especially when conducting experiments with particles, its effect on the organism strongly depends on its bioavailability, which can be modified by aggregation (Drago et al., 2020) and sedimentation (Schür et al., 2020). Until now, only a few studies used reference or natural particles in microplastic ecotoxicology tests with aquatic organisms. However, these and our results show that an urgent need exists to use a reference particle instead of a commonly used particle-free control to evaluate a potential plastic effect and thus a valid risk assessment of MP for aquatic organisms (Connors et al., 2017; Ogonowski et al., 2018).

### 4.3 High food vs. low food

In the AT<sub>24</sub> and AT<sub>96</sub>, we tested food limitation as an additional stressor, which was excluded in the CT, where sufficient food algae were available. After 24 h, food limitation exhibits no effect on survival, because *B. calyciflorus* is able to survive starvation during the first 24 h (Kirk, 1997). In the AT<sub>96</sub>, particle exposure caused comparable effects in the life history traits of *B. calyciflorus* for all three sizes and food/nonfood-particle proportions, and these differed markedly from the particle-free control. The most obvious cause for this is likely to be a dilution of the food alga (Kong and Koelmans, 2019). As a

consequence, the ingestion of food particles decreased, which resulted in declining energy uptake that ultimately affected life history traits (Arruda et al., 1983; McCabe and O'Brien, 1983). When comparing the results of the AT, this process can be seen by monitoring  $N(t)$ . After 24 h, no difference in survival was observed between particle-free (food algae only) and particle treatments (PS, Si), as sufficient food algae were available for the starting population of 7 rotifers. Only during the course (AT<sub>96</sub>) of the experiment and increasing population size, subsequently higher algae consumption rate potentially led to a deterioration of the food/nonfood-particle ratio, and the dilution effect became apparent. Further evidence for this explanation came from the life table experiments (CT), where the algae dilution factor (~6% of the diet) was low. In this case, enough food algae could be ingested to compensate the nutrient deficiency that occurred during the ingestion of nonfood particles. For this reason, the ratio between food/nonfood particles is crucial for life history trait outcomes, as food availability can alleviate nonfood particle (e.g. MPs) effects in *B. calyciflorus* (Drago and Weithoff, 2021; Xue et al., 2021) and *D. magna* (Ogonowski et al., 2016). Xue et al. (2021) reported a decrease in net reproduction rate during the exposure to polyethylene microbeads, but this effect could be mitigated by increasing the concentration of food algae. That supports our hypothesis that a diluting effect or the ratio between food/nonfood particles plays a more important role for life history traits of *B. calyciflorus* than a direct toxic effect of PS microplastics. In general, sufficient food sources often attenuate trait responses to toxic stress stimuli in rotifers (Cecchine and Snell, 1999; Sarma et al., 2001) and daphnids (Reinikainen et al., 1994; Pereira and Goncalves, 2007). Nevertheless, we found a decreased reproduction in the CT, even under high food conditions in the 3 μm treatment, which indicates that 3 μm MP was ingested along with the same sized algae (*M. minutum*: ~3.5 μm in diameter; Drago et al., 2020). In a study by DeMott (1986) with *B. calyciflorus*, researchers showed that no discrimination was found between particles with and without nutritional value; thus, *B. calyciflorus* does not seem to be able to discriminate between food algae and MP. In light of these findings, a risk of mistakenly ingestion of MPs appears when they are in the same size range as the corresponding food particles. Considering the many studies that demonstrated MP ingestion in zooplankton species and additionally the continuous release of plastic debris into aquatic ecosystems, as well as the continued degradation of plastics, researchers suggested that reproduction affected by MP ingestion could result in implications for entire food webs and associated ecosystem services.

### 4.4 Environmental relevance of the study

The concentrations of MP used in the experiment are, by far, higher than the ones that have been found in freshwater

ecosystems. However, higher MP concentrations compared to MP assessments in water columns of lakes and rivers are quite conceivable near point and nonpoint sources such as wastewater treatment plants (WWTP) or during/after storm or heavy rain events. For example, Hitchcock (2020) showed that the concentration of MP per m<sup>3</sup> water was increased 40-fold during a storm event. It should be noted that in most cases for MP concentration surveys, mesh widths of < 300 µm were used (Barrows et al., 2017; Mendoza and Balcer, 2019; Lindeque et al., 2020). However, this does not capture the fraction of small MP that falls within the food size spectrum of a broad range of zooplankton species (Cole et al., 2013; Botterell et al., 2019). The indication of underestimated high concentrations of small MP are shown in studies of Wiggin and Holland (2019) and Enders et al. (2015) for marine and estuarine ecosystems, where the smallest size fraction of MP (3–20 and 10 µm, resp.) accounted for the highest MP proportion. Thus, the expectation exists that the concentrations of small MP found in the future will increase by employing advanced methods and also due to continuous physical fragmentation of larger plastics (Andrady, 2011). However, it should be emphasized that we used pristine PS without additives like plasticizer, flame-retardants, or UV stabilizers, which are bioavailable for animals and are suspected to be toxic when ingested (Rochman, 2015). Microplastics are also known for their potential to adsorb organic pollutants like pesticides (Li et al., 2021), which could also be a threat for aquatic organisms. Their implementation might have altered our results, as shown in a study by Schrank et al. (2019) using *D. magna* and leaching additives.

## 5 Conclusion

We conclude that the combination of short- and long-term experiments in ecotoxicology provide a much deeper understanding of the underlying processes of potential toxicity. Further, this appears to be particularly true for sublethal effects such as delayed reproduction or changes in life span.

## Data availability statement

The original contributions presented in the study are included in the article/Supplementary Material, and further inquiries can be directed to the corresponding author.

## Author contributions

JP, DN, and GW designed the experiments. JP and DN conducted the experiment and performed the analysis of the data. JP, DN, CD, and GW wrote the manuscript. All authors discussed the results and provided extensive comments on the manuscript concerning the analysis, interpretation, and writing. All authors approved the final version and accepted accountability for all aspects of the work.

## Funding

This research was supported by the BMBF project MikroPlaTaS (02WPL1448C) granted to GW. Funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project number 491466077.

## Acknowledgments

We thank S. Donath, C. Schirmer, and S. Saumweber for providing us with samples and technical support. We also thank the reviewers for their comments that improved the manuscript.

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors, and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

## Supplementary material

The supplementary material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fenvs.2022.955425/full#supplementary-material>

## References

- Allen, S., Allen, D., Phoenix, V. R., Le Roux, G., Durántez Jiménez, P., Simonneau, A., et al. (2019). Atmospheric transport and deposition of microplastics in a remote mountain catchment. *Nat. Geosci.* 12, 339–344. doi:10.1038/s41561-019-0335-5
- An, D., Na, J., Song, J., and Jung, J. (2021). Size-dependent chronic toxicity of fragmented polyethylene microplastics to *Daphnia magna*. *Chemosphere* 271, 129591. doi:10.1016/j.chemosphere.2021.129591
- Ardrady, A. L. (2011). Microplastics in the marine environment. *Mar. Pollut. Bull.* 62, 1596–1605. doi:10.1016/j.marpolbul.2011.05.030
- Arndt, H. (1993). Rotifers as predators on components of the microbial web (bacteria, heterotrophic flagellates, ciliates) – a review. *Hydrobiologia* 255, 231–246. doi:10.1007/BF00025844
- Arruda, J. A., Marzholz, G. R., and Faulk, R. T. (1983). The role of suspended sediments in the nutrition of zooplankton in turbid reservoirs. *Ecology* 64, 1225–1235. doi:10.2307/1937831
- Barrows, A. P. W., Neumann, C. A., Bergera, M. L., and Shaw, S. D. (2017). Grab vs. neuston tow net: a microplastic sampling performance comparison and possible advances in the field. *Anal. Methods* 9, 1446–1453. doi:10.1039/c6ay02387h
- Begon, M., Harper, J. L., and Townsend, C. R. (1996). *Ecology: Individuals, populations, and communities*. 3rd edition. Cambridge, MA: Blackwell Science Ltd.
- Bergmann, M., Mützel, S., Primpke, S., Tekman, M. B., Trachsel, J., Gerdt, G., et al. (2019). White and wonderful? Microplastics prevail in snow from the alps to the arctic. *Sci. Adv.* 5, eaax1157. doi:10.1126/sciadv.aax1157
- Blettler, M. C. M., Abrial, E., Khan, F. R., Sivri, N., and Espinola, L. A. (2018). Freshwater plastic pollution: recognizing research biases and identifying knowledge gaps. *Water Res.* 143, 416–424. doi:10.1016/j.watres.2018.06.015
- Botterell, Z. L. R., Beaumont, N., Dorrington, T., Steinke, M., Thompson, R. C., Lindeque, P. K., et al. (2019). Bioavailability and effects of microplastics on marine zooplankton: a review. *Environ. Pollut.* 245, 98–110. doi:10.1016/j.envpol.2018.10.065
- Cappello, T., De Marco, G., Conti, G. O., Giannetto, A., Ferrante, M., Mauceri, A., et al. (2021). Time-dependent metabolic disorders induced by short-term exposure to polystyrene microplastics in the Mediterranean mussel *Mytilus galloprovincialis*. *Ecotoxicol. Environ. Saf.* 209, 111780. doi:10.1016/j.ecoenv.2020.111780
- Carpenter, E. J., and Smith, K. L., Jr. (1972). Plastics on the sargasso sea surface. *Science* 175, 1240–1241. doi:10.1126/science.175.4027.1240
- Castellanos Paez, M. E., Kurokura, H., and Kasahara, S. (1988). Embryonic development of amictic eggs of a rotifer *Brachionus plicatilis*. *J. Fac. Appl. Biol. Sci.* 27, 93–99.
- Cecchine, G., and Snell, T. W. (1999). Toxicant exposure increases threshold food levels in freshwater rotifer populations. *Environ. Toxicol.* 14, 523–530. doi:10.1002/(sici)1522-7278(199912)14:5<523::aid-tox6>3.0.co;2-2
- Coady, K. K., Burgoon, L., Doskey, C., and Davis, J. W. (2020). Assessment of transcriptomic and apical responses of *Daphnia magna* exposed to a polyethylene microplastic in a 21-d chronic study. *Environ. Toxicol. Chem.* 39, 1578–1589. doi:10.1002/etc.4745
- Cole, M., Lindeque, P., Fileman, E., Halsband, C., and Galloway, T. S. (2015). The impact of polystyrenemicroplastics on feeding, function and fecundity in the marine copepod *Calanus helgolandicus*. *Environ. Sci. Technol.* 49, 1130–1137. doi:10.1021/es504525u
- Cole, M., Lindeque, P., Fileman, E., Halsband, C., Goodhead, R., Moger, J., et al. (2013). Microplastic ingestion by zooplankton. *Environ. Sci. Technol.* 47, 6646–6655. doi:10.1021/es400663f
- Connors, K. A., Dyer, S. D., and Belanger, S. E. (2017). Advancing the quality of environmental microplastic research. *Environ. Toxicol. Chem.* 36, 1697–1703. doi:10.1002/etc.3829
- DeMott, W. R. (1986). The role of taste in food selection by freshwater zooplankton. *Oecologia* 69, 334–340. doi:10.1007/BF00377053
- Di, M., and Wang, J. (2018). Microplastics in surface waters and sediments of the three gorges reservoir, China. *Sci. Total Environ.* 616–617, 1620–1627. doi:10.1016/j.scitotenv.2017.10.150
- Drago, C., Pawlak, J., and Weithoff, G. (2020). Biogenic aggregation of small microplastics alters their ingestion by a common freshwater micro-invertebrate. *Front. Environ. Sci.* 8, 574274. doi:10.3389/fenvs.2020.574274
- Drago, C., and Weithoff, G. (2021). Variable fitness response of two rotifer species exposed to microplastics particles: the role of food quantity and quality. *Toxics* 9, 305. doi:10.3390/toxics9110305
- Eltens, Y. S., and Böhn, T. (2019). Acute and chronic effects of polystyrene microplastics on juvenile and adult *Daphnia magna*. *Environ. Pollut.* 254, 112919. doi:10.1016/j.envpol.2019.07.087
- Enders, K., Lenz, R., Stedmon, C. A., and Nielson, T. G. (2015). Abundance, size and polymer composition of marine microplastics  $\geq 10 \mu\text{m}$  in the Atlantic Ocean and their modelled vertical distribution. *Mar. Pollut. Bull.* 100, 70–81. doi:10.1016/j.marpolbul.2015.09.027
- Eriksen, M., Lebreton, L. C. M., Carson, H. S., Thiel, M., Moore, C. J., Borroero, J. C., et al. (2014). Plastic pollution in the world's oceans: more than 5 trillion plastic pieces weighing over 250,000 tons afloat at sea. *PLoS One* 9, e111913. doi:10.1371/journal.pone.0111913
- Fischer, E. K., Paglialonga, L., Czech, E., and Tamminga, M. (2016). Microplastic pollution in lakes and lake shoreline sediments – A case study on lake Bolsena and lake Chiusi (central Italy). *Environ. Pollut.* 213, 648–657. doi:10.1016/j.envpol.2016.03.012
- Fok, L., Cheng, N. Y. I., and Yeung, Y.-Y. (2019). “Mismanaged plastic waste: far side of the moon,” in *Environmental sustainability and education for waste management: implications for policy and practice*. Editors W. W. M. So, C. F. Chow, and J. C. K. Lee (Singapore: Springer), 57–71. doi:10.1007/978-981-13-9173-6\_5
- Frydkjær, C. K., Iversen, N., and Roslev, P. (2017). Ingestion and egestion of microplastics by the cladoceran *Daphnia magna*: effects of regular and irregular shaped plastic and sorbed phenanthrene. *Bull. Environ. Contam. Toxicol.* 99, 655–661. doi:10.1007/s00128-017-2186-3
- Fussmann, G. F., Weithoff, G., and Yoshida, T. (2005). A direct, experimental test of resource vs. consumer dependence. *Ecology* 86, 2924–2930. doi:10.1890/04-1107
- Gambardella, C., Morgana, S., Bramini, M., Rotini, A., Manfra, L., Migliore, L., et al. (2018). Ecotoxicological effects of polystyrene microbeads in a battery of marine organisms belonging to different trophic levels. *Mar. Environ. Res.* 141, 313–321. doi:10.1016/j.marenvres.2018.09.023
- Gambardella, C., Morgana, S., Ferrando, S., Bramini, M., Piazza, V., Costa, E., et al. (2017). Effects of polystyrene microbeads in marine planktonic crustaceans. *Ecotoxicol. Environ. Saf.* 145, 250–257. doi:10.1016/j.ecoenv.2017.07.036
- Guillard, R. R. L., and Lorenzen, C. J. (1972). Yellow-green algae with chlorophyllide C<sup>12</sup>. *J. Phycol.* 8, 10–14. doi:10.1111/j.1529-8817.1972.tb03995.x
- Harris, L. S. T., and Carrington, E. (2019). Impacts of microplastic vs. natural abiotic particles on the clearance rate of a marine mussel. *Limnol. Oceanogr. Lett.* 5, 66–73. doi:10.1002/lol2.10120
- Hart, R. C. (1988). Zooplankton feeding rates in relation to suspended sediment content: potential influences on community structure in a turbid reservoir. *Freshw. Biol.* 19, 123–139. doi:10.1111/j.1365-2427.1988.tb00334.x
- Higgins, J. P. T., Li, T., and Deeks, J. J. (2019). “Choosing effect measures and computing estimates of effect,” in *Cochrane handbook for systematic reviews of interventions*. Editors J. P. T. Higgins, J. Thomas, J. Chandler, M. Cumpston, T. Li, M. J. Page, et al. (Chichester: John Wiley and Sons) (Cochrane, 2019, version 6.0, updated July 2019).
- Hitchcock, J. N. (2020). Storm events as key moments of microplastic contamination in aquatic ecosystems. *Sci. Total Environ.* 734, 139436. doi:10.1016/j.scitotenv.2020.139436
- Imhof, H. K., Rusek, J., Thiel, M., Wolinska, J., and Laforsch, C. (2017). Do microplastic particles affect *Daphnia magna* at the morphological, life history and molecular level? *PLoS One* 12, e0187590. doi:10.1371/journal.pone.0187590
- Jaikumar, G., Brun, N. R., Vijver, M. G., and Bosker, T. (2019). Reproductive toxicity of primary and secondary microplastics to three cladocerans during chronic exposure. *Environ. Pollut.* 249, 638–646. doi:10.1016/j.envpol.2019.03.085
- Jeong, C.-B., Won, E.-J., Kang, H.-M., Lee, M.-C., Hwang, D.-S., Hwang, U.-K., et al. (2016). Microplastic size-dependent toxicity, oxidative stress induction, and p-JNK and p-p38 activation in the monogonont rotifer (*Brachionus koreanus*). *Environ. Sci. Technol.* 50, 8849–8857. doi:10.1021/acs.est.6b01441
- Kanhai, L. D. K., Officer, R., Lyashevskaya, O., Thompson, R. C., and O’Connor, I. (2017). Microplastic abundance, distribution and composition along a latitudinal gradient in the Atlantic Ocean. *Mar. Pollut. Bull.* 115, 307–314. doi:10.1016/j.marpolbul.2016.12.025
- Kapp, K. J., and Yeatman, E. (2018). Microplastic hotspots in the snake and lower columbia rivers: a journey from the greater yellowstone ecosystem to the pacific ocean. *Environ. Pollut.* 241, 1082–1090. doi:10.1016/j.envpol.2018.06.033
- Kirk, K. L., and Gilbert, J. J. (1990). Suspended clay and the population dynamics of planktonic rotifers and cladocerans. *Ecology* 71, 1741–1755. doi:10.2307/1937582
- Kirk, K. L. (1997). Life-history response to variable environments: starvation and reproduction in planktonic rotifers. *Ecology* 78, 434–441. doi:10.1890/0012-9658(1997)078[0434:LHRTVE]2.0.CO;2
- Kirk, L. K. (1991). Suspended clay reduces *Daphnia* feeding rate: behavioural mechanisms. *Freshw. Biol.* 25, 357–365. doi:10.1111/j.1365-2427.1991.tb00498.x

- Kong, X., and Koelmans, A. A. (2019). Modeling decreased resilience of shallow lake ecosystems toward eutrophication due to microplastic ingestion across the food web. *Environ. Sci. Technol.* 53, 13822–13831. doi:10.1021/acs.est.9b03905
- Lee, K.-W., Shim, W. J., Kwon, O. Y., and Kang, J.-H. (2013). Size-dependent effects of micro polystyrene particles in the marine copepod *Triptopus japonicas*. *Environ. Sci. Technol.* 47, 11278–11283. doi:10.1021/es401932b
- Li, H., Wang, F., Li, J., Deng, S., and Zhang, S. (2021). Adsorption of three pesticides on polyethylene microplastics in aqueous solutions: kinetics, isotherms, thermodynamics, and molecular dynamics simulation. *Chemosphere* 264, 128556. doi:10.1016/j.chemosphere.2020.128556
- Lindeque, P. K., Cole, M., Coppock, R. L., Lewis, C. N., Miller, R. Z., Watts, A. J. R., et al. (2020). Are we underestimating microplastic abundance in the marine environment? A comparison of microplastic capture with nets of different mesh-size. *Environ. Pollut.* 265, 114721. doi:10.1016/j.envpol.2020.114721
- Maes, T., Barry, J., Stenton, C., Roberts, E., Hicks, R., and Bignell, J. (2020). The world is your oyster: low-dose, long-term microplastic exposure of juvenile oysters. *Heliyon* 6, e03103. doi:10.1016/j.heliyon.2019.e03103
- Mani, T., Hauk, A., Walter, U., and Burkhardt-Holm, P. (2015). Microplastics profile along the rhine river. *Sci. Rep.* 5, 17988. doi:10.1038/srep17988
- McCabe, G. D., and O'Brien, W. J. (1983). The effects of suspended silt on feeding and reproduction of *Daphnia pulex*. *Am. Midl. Nat.* 110, 324. doi:10.2307/2425273
- Mendoza, L. M. R., and Balcer, M. (2019). Microplastics in freshwater environments: a review of quantification assessment. *TrAC Trends Anal. Chem.* 113, 402–408. doi:10.1016/j.trac.2018.10.020
- Naidu, S. A., Rao, V. R., and Ramu, K. (2018). Microplastics in the benthic invertebrates from the coastal waters of Kochi, southeastern Arabian Sea. *Environ. Geochem. Health* 40, 1377–1383. doi:10.1007/s10653-017-0062-z
- Ogonowski, M., Gerdes, Z., and Gorokhova, E. (2018). What we know and what we think we know about microplastic effects – a critical perspective. *Curr. Opin. Environ. Sci. Health* 1, 41–46. doi:10.1016/j.coesh.2017.09.001
- Ogonowski, M., Schür, C., Jarsen, A., and Gorokhova, E. (2016). The effects of natural and anthropogenic microparticles on individual fitness in *Daphnia magna*. *PLoS One* 11, e0155063. doi:10.1371/journal.pone.0155063
- Paraskevopoulou, S., Tiedemann, R., and Weithoff, G. (2018). Differential response to heat stress among evolutionary lineages of an aquatic invertebrate species complex. *Biol. Lett.* 14, 20180498. doi:10.1098/rsbl.2018.0498
- Pereira, J. L., and Goncalves, F. (2007). Effects of food availability on the acute and chronic toxicity of the insecticide methomyl to *Daphnia* spp. *Sci. Total Environ.* 386, 9–20. doi:10.1016/j.scitotenv.2007.07.040
- Phothakwanpracha, J., Lirdwitayaprasit, T., and Pairohakul, S. (2021). Effects of sizes and concentrations of different types of microplastics on bioaccumulation and lethality rate in the green mussel, *Perna viridis*. *Mar. Pollut. Bull.* 173, 112954. doi:10.1016/j.marpolbul.2021.112954
- Piehl, S., Leibner, A., Löder, M. G. J., Dris, R., Bogner, C., Laforsch, C., et al. (2018). Identification and quantification of macro- and microplastics on an agricultural farmland. *Sci. Rep.* 8, 17950. doi:10.1038/s41598-018-36172-y
- Qiao, R., Deng, Y., Zhang, S., Borri Wolosker, M., Zhu, Q., Ren, H., et al. (2019). Accumulation of different shapes of microplastics initiates intestinal injury and gut microbiota dysbiosis in the gut of zebrafish. *Chemosphere* 236, 124334. doi:10.1016/j.chemosphere.2019.07.065
- Reinikainen, M., Ketola, M., and Walls, M. (1994). Effects of the concentration of toxic *Microcystis aeruginosa* and alternative food on the survival of *Daphnia pulex*. *Limnol. Oceanogr.* 39, 424–432. doi:10.4319/lo.1994.39.2.0424
- Rochman, C. M. (2015). “The complex mixture, fate and toxicity of chemicals associated with plastic debris in the marine environment,” in *Marine Anthropogenic Litter*. Editors M. Bergmann, L. Gutow, and M. Klages (Cham: Springer). doi:10.1007/978-3-319-16510-3\_5
- Rothhaupt, K. O. (1990). Population growth rates of two closely related rotifer species: effects of food quantity, particle size, and nutritional quality. *Freshw. Biol.* 23, 561–570. doi:10.1111/j.1365-2427.1990.tb00295.x
- Sarma, S. S., Nandini, S., and Flores, J. L. (2001). Effect of methyl parathion on the population growth of the rotifer *Brachionus patulus* (O.F. Müller) under different algal food (*Chlorella vulgaris*) densities. *Ecotoxicol. Environ. Saf.* 48, 190–195. doi:10.1006/eesa.2000.2014
- Sarma, S. S., Ramírez Pérez, T., and Nandini, S. (2000). Comparison of the sensitivity of *Brachionus calyciflorus* and *Brachionus patulus* (Rotifera) to selected heavy metals under low and high food (*Chlorella vulgaris*) levels. *Bull. Environ. Contam. Toxicol.* 64, 735–739. doi:10.1007/s001280000065
- Schällicke, S., Sobisch, L.-Y., Martin-Creuzburg, D., and Wacker, A. (2019). Food quality–quantity co-limitation: interactive effects of dietary carbon and essential lipid supply on population growth of a freshwater rotifer. *Freshw. Biol.* 64, 903–912. doi:10.1111/fwb.13272
- Scheirer, C. J., Ray, W. S., and Hare, N. (1976). The analysis of ranked data derived from completely randomized factorial designs. *Biometrics* 32, 429. doi:10.2307/2529511
- Scherer, C., Brennholt, N., Reifferscheid, G., and Wagner, M. (2017). Feeding type and development drive the ingestion of microplastics by freshwater invertebrates. *Sci. Rep.* 7, 17006. doi:10.1038/s41598-017-17191-7
- Schrank, I., Trotter, B., Dummert, J., Scholz-Böttcher, B. M., Löder, M. G. J., Laforsch, C., et al. (2019). Effects of microplastic particles and leaching additive on the life history and morphology of *Daphnia magna*. *Environ. Pollut.* 255, 113233. doi:10.1016/j.envpol.2019.113233
- Schür, C., Zipp, S., Thalau, T., and Wagner, M. (2020). Microplastics but not natural particles induce multigenerational effects in *Daphnia magna*. *Environ. Pollut.* 260, 113904. doi:10.1016/j.envpol.2019.113904
- Setälä, O., Fleming-Lehtinen, V., and Lehtiniemi, M. (2014). Ingestion and transfer of microplastics in the planktonic food web. *Environ. Pollut.* 185, 77–83. doi:10.1016/j.envpol.2013.10.013
- Snell, T. W. (2014). Rotifers as models for the biology of aging. *Int. Rev. Hydrobiol.* 99, 84–95. doi:10.1002/iroh.201301707
- Stabel, H.-H. (1986). Calcite precipitation in Lake Constance: Chemical equilibrium, sedimentation, and nucleation by algae. *Limnol. Oceanogr.* 31, 1081–1094. doi:10.4319/lo.1986.31.5.1081
- Sun, Y., Xu, W., Gu, Q., Chen, Y., Zhou, Q., Zhang, L., et al. (2019). Small-sized microplastics negatively affect rotifers: changes in the key life-history traits and rotifer - *phaeocystis* population dynamics. *Environ. Sci. Technol.* 53, 9241–9251. doi:10.1021/acs.est.9b02893
- Tietjen, T., Vähätalo, A. V., and Wetzel, R. G. (2005). Effects of clay mineral turbidity on dissolved organic carbon and bacterial production. *Aquat. Sci.* 67, 51–60. doi:10.1007/s00027-004-0753-2
- Wagner, M., Scherer, C., Alvarez-Muñoz, D., Brennholt, N., Bourrain, X., Buchinger, S., et al. (2014). Microplastics in freshwater ecosystems: what we know and what we need to know. *Environ. Sci. Eur.* 26, 12. doi:10.1186/s12302-014-0012-7
- Wallace, R. L., and Snell, T. W. (2010). “Rotifera,” in *Ecology and classification of north American Freshwater invertebrates*. Editors J. H. Thorp and A. P. Covich (London: Academic Press 2010, Elsevier Inc.), 173–235. doi:10.1016/B978-0-12-374855-3.00008-X
- Walz, N. (1993). *Plankton regulation dynamics: Experiments and models in rotifer continuous cultures*. Heidelberg: Springer-Verlag Berlin.
- Wang, Y., Zhang, D., Zhang, M., Mu, J., Ding, G., Mao, Z., et al. (2019). Effects of ingested polystyrene microplastics on brine shrimp, *Artemia parthenogenetica*. *Environ. Pollut.* 244, 715–722. doi:10.1016/j.envpol.2018.10.024
- Weithoff, G. (2007). Dietary restriction in two rotifer species: the effect of the length of food deprivation on life span and reproduction. *Oecologia* 153, 303–308. doi:10.1007/s00442-007-0739-6
- Weithoff, G., and Wacker, A. (2007). The mode of nutrition of mixotrophic flagellates determines the food quality for their consumers. *Funct. Ecol.* 21, 1092–1098. doi:10.1111/j.1365-2435.2007.01333.x
- Weithoff, G., and Walz, N. (1995). Influence of filamentous cyanobacterium *Planktothrix agardhii* on population growth and reproductive pattern of the rotifer *Brachionus calyciflorus*. *Hydrobiologia* 313, 381–386. doi:10.1007/BF00025974
- Wiggin, K. J., and Holland, E. B. (2019). Validation and application of cost and time effective methods for the detection of 3–500 µm sized microplastics in the urban marine and estuarine environments surrounding Long Beach, California. *Mar. Pollut. Bull.* 143, 152–162. doi:10.1016/j.marpolbul.2019.03.060
- Wilkinson, G. M., Carpenter, S. R., Cole, J. J., Pace, M. L., and Yang, C. (2013). Terrestrial support of pelagic consumers: patterns and variability revealed by a multilake study. *Freshw. Biol.* 58, 2037–2049. doi:10.1111/fwb.12189
- Windsor, F. M., Tilley, R. M., Tyler, C. R., and Ormerod, S. J. (2019). Microplastic ingestion by riverine macroinvertebrates. *Sci. Total Environ.* 646, 68–74. doi:10.1016/j.scitotenv.2018.07.271
- Xue, Y.-H., Sun, Z.-X., Feng, L.-S., Jin, T., Xing, J.-C., Wen, X.-L., et al. (2021). Algal density affects the influences of polyethylene microplastics on the freshwater rotifer *Brachionus calyciflorus*. *Chemosphere* 270, 128613. doi:10.1016/j.chemosphere.2020.128613
- Yap, V. H. S., Chase, Z., Wright, J. T., Hurd, C. L., Lavers, J. L., Lenz, M., et al. (2020). A comparison with natural particles reveals a small specific effect of PVC microplastics on mussel performance. *Mar. Pollut. Bull.* 160, 111703. doi:10.1016/j.marpolbul.2020.111703
- Zimmermann, L., Göttlich, S., Oehlmann, J., Wagner, M., and Völker, C. (2020). What are the drivers of microplastic toxicity? Comparing the toxicity of plastic chemicals and particles to *Daphnia magna*. *Environ. Pollut.* 267, 115392. doi:10.1016/j.envpol.2020.115392